

[Go to Product page](#)

Datasheet for ABIN1698920

**anti-Supervillin antibody (AA 51-150) (Alexa Fluor 647)**

## Overview

|                      |  |
|----------------------|--|
| Quantity:            | 100 µL   |
| Target:              | Supervillin (SVIL)   |
| Binding Specificity: | AA 51-150  |
| Reactivity:          | Mouse  |
| Host:                | Rabbit   |
| Clonality:           | Polyclonal   |
| Conjugate:           | This Supervillin antibody is conjugated to Alexa Fluor 647   |
| Application:         | Western Blotting (WB), Immunofluorescence (Cultured Cells) (IF (cc)), Immunofluorescence (Paraffin-embedded Sections) (IF (p)) |

## Product Details

|                       |   |
|-----------------------|---|
| Immunogen:            | KLH conjugated synthetic peptide derived from human Supervillin |
| Isotype:              | IgG   |
| Cross-Reactivity:     | Mouse   |
| Predicted Reactivity: | Human,Rat,Dog,Cow,Sheep,Pig,Horse,Chicken                       |
| Purification:         | Purified by Protein A.  |

## Target Details

|                   |   |
|-------------------|---|
| Target:           | Supervillin (SVIL)                            |
| Alternative Name: | Supervillin ( <a href="#">SVIL Products</a> ) |

## Target Details

|             |  |
|-------------|--|
| Background: | <p>Synonyms: Supervillin, Archvillin, p205/p250, SVIL</p> <p>Background: Forms a high-affinity link between the actin cytoskeleton and the membrane. Isoform 1 (archvillin) is among the first costameric proteins to assemble during myogenesis and it contributes to myogenic membrane structure and differentiation. Appears to be involved in myosin II assembly. May modulate myosin II regulation through MLCK during cell spreading, an initial step in cell migration. May play a role in invadopodial function. Isoform 2 may be involved in modulation of focal adhesions. Supervillin-mediated down-regulation of focal adhesions involves binding to TRIP6. Plays a role in cytokinesis through KIF14 interaction (By similarity).</p> |
|-------------|--|

|          |      |
|----------|------|
| Gene ID: | 6840 |
|----------|------|

|          |                        |
|----------|------------------------|
| UniProt: | <a href="#">O95425</a> |
|----------|------------------------|

## Application Details

|                    |  |
|--------------------|--|
| Application Notes: | IF(IHC-P) 1:50-200<br>IF(IHC-F) 1:50-200<br>IF(ICC) 1:50-200 |
|--------------------|--|

|               |                       |
|---------------|-----------------------|
| Restrictions: | For Research Use only |
|---------------|-----------------------|

## Handling

|                    |  |
|--------------------|--|
| Format:            | Liquid   |
| Concentration:     | 1 µg/µL  |
| Buffer:            | Aqueous buffered solution containing 0.01M TBS ( pH 7.4) with 1 % BSA, 0.03 % Proclin300 and 50 % Glycerol.        |
| Preservative:      | ProClin  |
| Precaution of Use: | This product contains ProClin: a POISONOUS AND HAZARDOUS SUBSTANCE, which should be handled by trained staff only. |
| Storage:           | -20 °C   |
| Storage Comment:   | Store at -20°C. Aliquot into multiple vials to avoid repeated freeze-thaw cycles.                                  |
| Expiry Date:       | 12 months  |