

Datasheet for ABIN1698930

anti-Ataxin 1 antibody (pSer775) (AbBy Fluor® 647)[Go to Product page](#)

Overview

Quantity:	100 µL
Target:	Ataxin 1 (ATXN1)
Binding Specificity:	pSer775
Reactivity:	Human, Mouse, Rat
Host:	Rabbit
Clonality:	Polyclonal
Conjugate:	This Ataxin 1 antibody is conjugated to AbBy Fluor® 647
Application:	Western Blotting (WB), Immunofluorescence (Cultured Cells) (IF (cc)), Immunofluorescence (Paraffin-embedded Sections) (IF (p))

Product Details

Immunogen:	KLH conjugated synthetic phosphopeptide derived from human Ataxin 1 around the phosphorylation site of Ser776
Isotype:	IgG
Cross-Reactivity:	Human, Mouse, Rat
Predicted Reactivity:	Dog,Cow,Sheep,Pig,Horse,Chicken,Rabbit
Purification:	Purified by Protein A.

Target Details

Target:	Ataxin 1 (ATXN1)
---------	------------------

Target Details

Alternative Name:	Ataxin 1 (ATXN1 Products)
Background:	Synonyms: Ataxin 1 phospho S776,p-Ataxin 1 phospho S776, ATX1, ATXN1, SCA1, Ataxin 1, Ataxin-1, Ataxin1, Spinocerebellar ataxia type 1, ATX1_HUMAN. Background: Ataxin-1, also designated spinocerebellar ataxia type 1 protein (Sca-1), is differentially expressed and localizes to both the cytoplasm and the nucleus. Mutations in Ataxin-1 are associated with the onset of the autosomal dominant neurodegenerative disorder spinocerebellar ataxia type 1 (SCA-1), which is characterized by progressive neuronal loss in the cerebellum, muscle wasting and ataxia. In Purkinje cells, where SCA-1 is predominantly observed, Ataxin-1 has been shown to directly associate with the Purkinje-enriched leucine-rich acidic nuclear protein (LANP) and the nuclear matrix-associated protein promyelocytic leukemia protein PML. In SCA-1, Ataxin-1 is mutated to encode a polyglutamine protein that forms nuclear aggregates, which interact significantly more strongly with LANP and contribute to the pathogenesis of SCA-1.
Gene ID:	6310
Pathways:	Synaptic Membrane

Application Details

Application Notes:	IF(IHC-P) 1:50-200 IF(IHC-F) 1:50-200 IF(ICC) 1:50-200
Restrictions:	For Research Use only

Handling

Format:	Liquid
Concentration:	1 µg/µL
Buffer:	Aqueous buffered solution containing 0.01M TBS (pH 7.4) with 1 % BSA, 0.03 % Proclin300 and 50 % Glycerol.
Preservative:	ProClin
Precaution of Use:	This product contains ProClin: a POISONOUS AND HAZARDOUS SUBSTANCE, which should be handled by trained staff only.
Storage:	-20 °C

Handling

Storage Comment:	Store at -20°C. Aliquot into multiple vials to avoid repeated freeze-thaw cycles.
Expiry Date:	12 months