

[Go to Product page](#)

Datasheet for ABIN1699287

**anti-C6orf130 antibody (AA 51-150) (Alexa Fluor 647)**

## Overview

Quantity:	100 µL
Target:	C6orf130 (OARD1)
Binding Specificity:	AA 51-150
Reactivity:	Human
Host:	Rabbit
Clonality:	Polyclonal
Conjugate:	This C6orf130 antibody is conjugated to Alexa Fluor 647
Application:	Western Blotting (WB), Immunofluorescence (Cultured Cells) (IF (cc)), Immunofluorescence (Paraffin-embedded Sections) (IF (p))

## Product Details

Immunogen:	KLH conjugated synthetic peptide derived from human C6orf130
Isotype:	IgG
Predicted Reactivity:	Human, Mouse, Rat, Dog, Pig, Horse, Rabbit
Purification:	Purified by Protein A.

## Target Details

Target:	C6orf130 (OARD1)
Alternative Name:	C6orf130 ( <a href="#">OARD1 Products</a> )
Background:	Synonyms: C6orf130, OARD1_HUMAN, Chromosome 6 open reading frame 130, dJ34B21.3, O

## Target Details

acetyl ADP ribose deacetylase C6orf130, Uncharacterized protein C6orf130.

Background: C6orf130 is a Making up nearly 6 % of the human genome, chromosome 6 contains around 1,200 genes within 170 million base pairs of sequence. Deletion of a portion of the q arm of chromosome 6 is associated with early onset intestinal cancer suggesting the presence of a cancer susceptibility locus. Porphyria cutanea tarda is associated with chromosome 6 through the HFE gene which, when mutated, predisposes an individual to developing this porphyria. Notably, the PARK2 gene, which is associated with Parkinson's disease, and the genes encoding the major histocompatibility complex proteins, which are key molecular components of the immune system and determine predisposition to rheumatic diseases, are also located on chromosome 6. Stickler syndrome, 21-hydroxylase deficiency and maple syrup urine disease are also associated with genes on chromosome 6. A bipolar disorder susceptibility locus has been identified on the q arm of chromosome 6. The C6orf130 gene product has been provisionally designated C6orf130 pending further characterization.

Gene ID: 221443

## Application Details

Application Notes: IF(IHC-P) 1:50-200  
IF(IHC-F) 1:50-200  
IF(ICC) 1:50-200

Restrictions: For Research Use only

## Handling

Format: Liquid

Concentration: 1 µg/µL

Buffer: Aqueous buffered solution containing 0.01M TBS ( pH 7.4) with 1 % BSA, 0.03 % Proclin300 and 50 % Glycerol.

Preservative: ProClin

Precaution of Use: This product contains ProClin: a POISONOUS AND HAZARDOUS SUBSTANCE, which should be handled by trained staff only.

Storage: -20 °C

Storage Comment: Store at -20°C. Aliquot into multiple vials to avoid repeated freeze-thaw cycles.

Expiry Date: 12 months