# antibodies -online.com





# anti-NCAPG antibody (AA 161-170) (Alexa Fluor 647)



| $\sim$ |     |      |     |
|--------|-----|------|-----|
|        | N/P | r\/I | i⊢₩ |

| Quantity:            | 100 μL   |
|----------------------|--|
| Target:              | NCAPG  |
| Binding Specificity: | AA 161-170   |
| Reactivity:          | Human  |
| Host:                | Rabbit   |
| Clonality:           | Polyclonal   |
| Conjugate:           | This NCAPG antibody is conjugated to Alexa Fluor 647   |
| Application:         | Western Blotting (WB), Immunofluorescence (Cultured Cells) (IF (cc)), Immunofluorescence (Paraffin-embedded Sections) (IF (p)) |

#### **Product Details**

| Immunogen:            | KLH conjugated synthetic peptide derived from human HCAP G |
|-----------------------|--|
| Isotype:              | IgG  |
| Cross-Reactivity:     | Human  |
| Predicted Reactivity: | Mouse,Rat,Dog,Pig,Horse,Rabbit                             |
| Purification:         | Purified by Protein A.                                     |

## **Target Details**

| Target:           | NCAPG                   |
|-------------------|-------------------------|
| Alternative Name: | Hcap G (NCAPG Products) |

## **Target Details**

| Background:         | Synonyms: CAPG, CHCG, Chromosome associated protein G, Chromosome condensation                   |  |
|---------------------|--|--|
|                     | protein G, Condensin complex subunit 3, Condensin subunit CAP G, HCAP G, Melanoma antiger        |  |
|                     | NY MEL 3, NCAPG, Non SMC condensin I complex, subunit G, NY MEL 3, XCAP G homolog,               |  |
|                     | CND3_HUMAN.  |  |
|                     | Background: HCAP G is a subunit of the condensin complex, which is responsible for the           |  |
|                     | condensation and stabilization of chromosomes during mitosis and meiosis. Phosphorylation        |  |
|                     | of the encoded protein activates the condensin complex. There are pseudogenes for this gene      |  |
|                     | on chromosomes 8 and 15. Alternative splicing results in multiple transcript variants. [provided |  |
|                     | by RefSeq, Aug 2012].  |  |
| Gene ID:            | 64151  |  |
| Application Details |  |  |
| Application Notes:  | IF(IHC-P) 1:50-200   |  |
|                     | IF(IHC-F) 1:50-200   |  |
|                     | IF(ICC) 1:50-200   |  |
| Restrictions:       | For Research Use only  |  |
| Handling            |  |  |
| Format:             | Liquid   |  |
| Concentration:      | 1 μg/μL  |  |
| Buffer:             | Aqueous buffered solution containing 0.01M TBS (pH 7.4) with 1 % BSA, 0.03 % Proclin300 and      |  |
|                     | 50 % Glycerol.   |  |
| Preservative:       | ProClin  |  |
| Precaution of Use:  | This product contains ProClin: a POISONOUS AND HAZARDOUS SUBSTANCE, which should be              |  |
|                     | handled by trained staff only.   |  |
| Storage:            | -20 °C   |  |
| Storage Comment:    | Store at -20°C. Aliquot into multiple vials to avoid repeated freeze-thaw cycles.                |  |
| Expiry Date:        | 12 months  |  |
|                     |  |  |