antibodies -online.com







anti-CSRP2BP antibody (AA 71-170) (AbBy Fluor® 647)



()	11/0	K\ /	iew	1
	\cup	'I V/I	$\square \vee \vee$	ı

Quantity:	100 μL	
Target:	CSRP2BP	
Binding Specificity:	AA 71-170	
Reactivity:	Human	
Host:	Rabbit	
Clonality:	Polyclonal	
Conjugate:	This CSRP2BP antibody is conjugated to AbBy Fluor® 647	
Application:	Western Blotting (WB), Immunofluorescence (Cultured Cells) (IF (cc)), Immunofluorescence (Paraffin-embedded Sections) (IF (p))	

Product Details

Immunogen:	KLH conjugated synthetic peptide derived from human CSRP2BP
Isotype:	IgG
Predicted Reactivity: Human, Mouse, Rat, Dog, Cow, Sheep, Pig, Horse, Rabbit	
Purification:	Purified by Protein A.

Target Details

Target:	CSRP2BP	
Alternative Name:	Name: CSRP2BP (CSRP2BP Products)	
Synonyms: ADA2A containing complex subunit 2, ADA2A-containing com		

CRP2 binding partner, CRP2-binding partner, CRP2BP, CSR2B_HUMAN, CSRP2 binding protein, CSRP2-binding protein, Csrp2bp, Cysteine rich protein 2 binding protein, Cysteine-rich protein 2-binding protein.

Background: CRP2BP is a 782 amino acid protein encoded by the human gene CSRP2BP. CRP2BP specifically interacts with the double LIM domain protein CRP2. The LIM domain is a conserved cysteine and histidine-containing structural module of two tandemly arranged zinc fingers. It has been identified in single or multiple copies in a variety of regulatory proteins, either in combination with defined functional domains, like homeodomains, or alone, like in the CRP family of LIM proteins. Members of the cysteine- and glycine-rich protein family (CRP1, CRP2 and CRP3) contain two zinc-binding LIM domains, LIM1 (amino-terminal) and LIM2 (carboxyl-terminal), and are implicated in diverse cellular processes linked to differentiation, growth control and pathogenesis. Although present in cytoplasm, CRP2BP is mainly a ubiquitously expressed nuclear protein, with highest expression in skeletal muscle and heart.

Gene ID:

57325

Application Details

Λnn	lication	Notes.	

IF(IHC-P) 1:50-200

IF(IHC-F) 1:50-200

IF(ICC) 1:50-200

Restrictions:

For Research Use only

Handling

Format:	Liquid	
Concentration:	1 μg/μL	
Buffer:	Aqueous buffered solution containing 0.01M TBS (pH 7.4) with 1 % BSA, 0.03 % Proclin300 and 50 % Glycerol.	
Preservative:	ProClin	
Precaution of Use:	This product contains ProClin: a POISONOUS AND HAZARDOUS SUBSTANCE, which should be handled by trained staff only.	
Storage:	-20 °C	
Storage Comment:	Store at -20°C. Aliquot into multiple vials to avoid repeated freeze-thaw cycles.	
Expiry Date:	12 months	