# antibodies .- online.com





# anti-DHX15 antibody (AA 101-200) (Alexa Fluor 647)



Go to Product page

( )	11/0	K\ /	iew	1
	$\cup$	ועוי	$\square \vee \vee$	ı

Quantity:	100 μL	
Target:	DHX15	
Binding Specificity:	AA 101-200	
Reactivity:	Human	
Host:	Rabbit	
Clonality:	Polyclonal	
Conjugate:	This DHX15 antibody is conjugated to Alexa Fluor 647	
Application:	Western Blotting (WB), Immunofluorescence (Cultured Cells) (IF (cc)), Immunofluorescence (Paraffin-embedded Sections) (IF (p))	

## **Product Details**

Immunogen:	KLH conjugated synthetic peptide derived from human DHX15	
Isotype:	IgG	
Predicted Reactivity: Human, Mouse, Rat, Dog, Cow, Pig, Horse, Rabbit		
Purification:	Purified by Protein A.	

# **Target Details**

Target:	DHX15	
Alternative Name:	DHX15 (DHX15 Products)	
Synonyms: ATP dependent RNA helicase #46, DBP1, DDX15, DEADH box 15, DEAH Asp Glu		

His box polypeptide 15, DEAH box polypeptide 15, DEAH box protein 15, HRH2, PRP43, PrPp43p, RNA, DHX15\_HUMAN.

Background: DEAD-box proteins, characterized by the conserved motif Asp-Glu-Ala-Asp, are putative RNA helicases implicated in several cellular processes involving modifications of RNA secondary structure and ribosome/spliceosome assembly. Based on their distribution patterns, some members of this family may be involved in embryogenesis, spermatogenesis, and cellular growth and division. DDX15 (DEAH-box protein 15), also known as DHX15, DBP1 or HRH2, is a nuclear ATP-dependent RNA helicase that is involved in pre-mRNA splicing and is a member of the DEAH-box subfamily of DEAD-box proteins. Expressed throughout the body, DDX15 is a pre-mRNA processing factor that plays a role in spliceosome disassembly after the release of mature mRNA. When localized to the nucleoli, DDX15 is thought to interact with the La/SSB autoantigen, an RNA chaperone that functions in various intracellular processes. DDX15 is 795 amino acids in length and is the human ortholog of the S. cerevisiae protein Prp43. helicase 2.

Gene ID:

1665

### **Application Details**

<b>Application Notes:</b>	

IF(IHC-P) 1:50-200

IF(IHC-F) 1:50-200

IF(ICC) 1:50-200

Restrictions:

For Research Use only

#### Handling

Format:	Liquid	
Concentration:	1 μg/μL	
Buffer:	Aqueous buffered solution containing 0.01M TBS (pH 7.4) with 1 % BSA, 0.03 % Proclin300 and 50 % Glycerol.	
Preservative:	ProClin	
Precaution of Use:	This product contains ProClin: a POISONOUS AND HAZARDOUS SUBSTANCE, which should be handled by trained staff only.	
Storage:	-20 °C	
Storage Comment:	Store at -20°C. Aliquot into multiple vials to avoid repeated freeze-thaw cycles.	
Expiry Date:	12 months	