

[Go to Product page](#)

Datasheet for ABIN1699430

anti-EXOSC10 antibody (AA 41-140) (Alexa Fluor 647)

Overview

| | |
|----------------------|--|
| Quantity: | 100 µL |
| Target: | EXOSC10 |
| Binding Specificity: | AA 41-140 |
| Reactivity: | Human, Mouse |
| Host: | Rabbit |
| Clonality: | Polyclonal |
| Conjugate: | This EXOSC10 antibody is conjugated to Alexa Fluor 647 |
| Application: | Western Blotting (WB), Immunofluorescence (Cultured Cells) (IF (cc)), Immunofluorescence (Paraffin-embedded Sections) (IF (p)) |

Product Details

| | |
|-----------------------|--|
| Immunogen: | KLH conjugated synthetic peptide derived from human EXOSC10/PMSCL2 |
| Isotype: | IgG |
| Cross-Reactivity: | Human, Mouse |
| Predicted Reactivity: | Rat,Dog,Cow,Sheep,Pig,Horse |
| Purification: | Purified by Protein A. |

Target Details

| | |
|-------------------|---|
| Target: | EXOSC10 |
| Alternative Name: | EXOSC10/PMSCL2 (EXOSC10 Products) |

Target Details

| | |
|-------------|---|
| Background: | <p>Synonyms: Autoantigen PM/Scl 2, Exosc10, Exosome component 10, EXOSX_HUMAN, P100 polymyositis scleroderma overlap syndrome associated autoantigen, P100 polymyositis-scleroderma overlap syndrome-associated autoantigen, p2, p3, p4, PM Scl, PM/Scl 100, PM/Scl-100, PMSCL, PMSCL2, Polymyositis/scleroderma autoantigen 100 kDa, Polymyositis/scleroderma autoantigen 2 100 kDa, Polymyositis/scleroderma autoantigen 2, RRP6, Rrp6p.</p> <p>Background: The exosome is a multi-subunit complex composed of several highly conserved proteins, some of which are 3' to 5' exonucleases. The complex is involved in a variety of cellular processes and is responsible for degrading unstable mRNAs that contain AU-rich (ARE) elements in their untranslated 3' region. EXOSC10, also known as PMSCL, PMSCL2, p2, p3, p4, RRP6, Rrp6p, PM-Scl, or PM/Scl-100, is an 885 amino acid protein that contains one HRDC domain and one 3'-5' exonuclease domain. Localized to both the cytoplasm and the nucleus, EXOSC10 is part of the post-splicing exosome complex and is involved in mRNA surveillance, mRNA nuclear export and nonsense-mediated decay of mRNAs containing premature stop codons. against EXOSC10 have been found in patients with scleroderma and/or polymyositis (chronic diseases of the skin and muscle, respectively), suggesting that EXOSC10 may be involved in the pathogenesis of these diseases. Two isoforms of EXOSC10 exist due to alternative splicing events.</p> |
|-------------|---|

| | |
|----------|------|
| Gene ID: | 5394 |
|----------|------|

Application Details

| | |
|--------------------|--|
| Application Notes: | IF(IHC-P) 1:50-200 IF(IHC-F) 1:50-200 IF(ICC) 1:50-200 |
|--------------------|--|

| | |
|---------------|-----------------------|
| Restrictions: | For Research Use only |
|---------------|-----------------------|

Handling

| | |
|--------------------|---|
| Format: | Liquid |
| Concentration: | 1 µg/µL |
| Buffer: | Aqueous buffered solution containing 0.01M TBS (pH 7.4) with 1 % BSA, 0.03 % Proclin300 and 50 % Glycerol. |
| Preservative: | ProClin |
| Precaution of Use: | This product contains ProClin: a POISONOUS AND HAZARDOUS SUBSTANCE, which should be |

Handling

| | |
|------------------|---|
| | handled by trained staff only. |
| Storage: | -20 °C |
| Storage Comment: | Store at -20°C. Aliquot into multiple vials to avoid repeated freeze-thaw cycles. |
| Expiry Date: | 12 months |