# antibodies - online.com







## anti-GLI4 antibody (AA 271-376) (AbBy Fluor® 647)



( )	ve	K\ /		A .
	$\cup$	1 V/	-	V۷

Quantity:	100 μL
Target:	GLI4
Binding Specificity:	AA 271-376
Reactivity:	Human
Host:	Rabbit
Clonality:	Polyclonal
Conjugate:	This GLI4 antibody is conjugated to AbBy Fluor® 647
Application:	Western Blotting (WB), Immunofluorescence (Cultured Cells) (IF (cc)), Immunofluorescence (Paraffin-embedded Sections) (IF (p))

## **Product Details**

Immunogen:	KLH conjugated synthetic peptide derived from human GLI4
Isotype:	IgG
Cross-Reactivity:	Human
Predicted Reactivity:	Mouse,Rat,Cow,Pig,Horse
Purification:	Purified by Protein A.

## **Target Details**

Target:	GLI4
Alternative Name:	GLI4 (GLI4 Products)

## Target Details

Racko	round:
Dacke	n Our iu.

Synonyms: GLI 4, GLI Kruppel family member GLI 4, GLI Kruppel family member GLI4, GLI4, GLI4, GLI4, GLI4, GLI4, HUMAN, HKR 4, HKR4, HKR4 protein, Krueppel related zinc finger protein 4, Krueppel-related zinc finger protein 4, Neural specic DNA binding protein, Oncogene HKR 4, Oncogene HKR4, Protein HKR 4, Protein HKR4, Zinc finger protein GLI 4, Zinc finger protein GLI4. Background: Zinc-finger proteins contain DNA-binding domains and have a wide variety of functions, most of which encompass some form of transcriptional activation or repression. The majority of zinc-finger proteins contain a krueppel-type DNA binding domain and a KRAB domain, which is thought to interact with KAP1, thereby recruiting histone modifying proteins. GLI-4, also known as HKR4, is a 376 amino acid protein that localizes to the nucleus and contains seven C2H2-type zinc fingers. Belonging to the krueppel C2H2-type zinc-finger protein family, GLI-4 may function as a transcriptional regulator, effectively activating or repressing the transcription of target genes. The gene encoding GLI-4 maps to human chromosome 8, which consists of nearly 146 million base pairs, houses more than 800 genes and is associated with a variety of diseases and malignancies.

Gene ID:

2738

### **Application Details**

Ann	lication	Notes

IF(IHC-P) 1:50-200

IF(IHC-F) 1:50-200

IF(ICC) 1:50-200

Restrictions:

For Research Use only

#### Handling

Format:	Liquid
Concentration:	1 μg/μL
Buffer:	Aqueous buffered solution containing 0.01M TBS (pH 7.4) with 1 % BSA, 0.03 % Proclin300 and 50 % Glycerol.
Preservative:	ProClin
Precaution of Use:	This product contains ProClin: a POISONOUS AND HAZARDOUS SUBSTANCE, which should be handled by trained staff only.
Storage:	-20 °C
Storage Comment:	Store at -20°C. Aliquot into multiple vials to avoid repeated freeze-thaw cycles.

		1.	
$\vdash$	land	lın	
	ıaııu	1111	U

Expiry Date:

12 months