# antibodies -online.com





# anti-C9ORF62 antibody (AA 51-150) (Biotin)



Go to Product page

| ( ) | 11/               | IN   | /ie    | A .    |
|-----|-------------------|------|--------|--------|
|     | / // <del> </del> | ۱ ات | /   (− | ' \/\/ |
|     |                   |      |        |        |

| Quantity:            | 100 μL  |  |
|----------------------|---|--|
| Target:              | C9ORF62   |  |
| Binding Specificity: | AA 51-150   |  |
| Reactivity:          | Human   |  |
| Host:                | Rabbit  |  |
| Clonality:           | Polyclonal  |  |
| Conjugate:           | This C9ORF62 antibody is conjugated to Biotin   |  |
| Application:         | Western Blotting (WB), Immunohistochemistry (Frozen Sections) (IHC (fro)), Immunohistochemistry (Paraffin-embedded Sections) (IHC (p)), ELISA |  |
|                      |   |  |

## **Product Details**

| Immunogen:            | KLH conjugated synthetic peptide derived from human C9orf62 |  |
|-----------------------|---|--|
| Isotype:              | IgG   |  |
| Predicted Reactivity: | Human   |  |
| Purification:         | Purified by Protein A.                                      |  |

# **Target Details**

| Target:           | C9ORF62  |  |
|-------------------|--|--|
| Alternative Name: | C9orf62 (C9ORF62 Products)   |  |
| Background:       | Synonyms: Chromosome 9 open reading frame 62, MGC35463, Putative uncharacterized |  |

protein C9orf62,CI062\_HUMAN.

Background: Chromosome 9 consists of about 145 million bases and 4 % of the human genome and encodes nearly 900 genes. Considered to play a role in gender determination, deletion of the distal portion of 9p can lead to development of male to female sex reversal, the phenotype of a female with a male X,Y genotype. Hereditary hemorrhagic telangiectasia, which is characterized by harmful vascular defects, is associated with the chromosome 9 gene encoding endoglin protein, ENG. Familial dysautonomia is also associated with chromosome 9 though through the gene IKBKAP. Notably, chromosome 9 encompasses the largest interferon family gene cluster. Chromosome 9 is partnered with chromosome 22 in the translocation leading to the aberrant production of BCR-ABL fusion protein often found in leukemias. The C9orf62 gene product has been provisionally designated C9orf62 pending further characterization.

Gene ID:

157927

### **Application Details**

Application Notes: WB 1:300-5000

IHC-P 1:200-400

IHC-F 1:100-500

Restrictions: For Research Use only

#### Handling

| Format:            | Liquid   |  |
|--------------------|--|--|
| Concentration:     | 1 μg/μL  |  |
| Buffer:            | Aqueous buffered solution containing 0.01M TBS (pH 7.4) with 1 % BSA, 0.03 % Proclin300 and 50 % Glycerol.         |  |
| Preservative:      | ProClin  |  |
| Precaution of Use: | This product contains ProClin: a POISONOUS AND HAZARDOUS SUBSTANCE, which should be handled by trained staff only. |  |
| Storage:           | -20 °C   |  |
| Storage Comment:   | Store at -20°C for 12 months.  |  |
| Expiry Date:       | 12 months  |  |