

[Go to Product page](#)

Datasheet for ABIN1700068

anti-RGS14 antibody (AA 151-250) (Biotin)

Overview

Quantity:	100 µL
Target:	RGS14
Binding Specificity:	AA 151-250
Reactivity:	Mouse, Rat
Host:	Rabbit
Clonality:	Polyclonal
Conjugate:	This RGS14 antibody is conjugated to Biotin
Application:	Western Blotting (WB), ELISA, Immunohistochemistry (Paraffin-embedded Sections) (IHC (p)), Immunohistochemistry (Frozen Sections) (IHC (fro))

Product Details

Immunogen:	KLH conjugated synthetic peptide derived from human RGS14
Isotype:	IgG
Cross-Reactivity:	Mouse, Rat
Predicted Reactivity:	Human,Cow
Purification:	Purified by Protein A.

Target Details

Target:	RGS14
Alternative Name:	RGS14 (RGS14 Products)

Target Details

Background:	<p>Synonyms: Highly similar to rap1/rap2 interacting protein, OTTHUMP00000223586, OTTHUMP00000223587, Regulation of G protein signaling 14, Regulator of G protein signaling 14, Regulator of G protein signalling 14, Regulator of G-protein signaling 14, RGS 14, RGS14, RGS14_HUMAN.</p> <p>Background: RGS14 is a member of the regulator of G-protein signaling family. This protein contains one RGS domain, two Raf-like Ras-binding domains (RBDs), and one GoLoco domain. The protein attenuates the signaling activity of G-proteins by binding, through its GoLoco domain, to specific types of activated, GTP-bound G alpha subunits. Acting as a GTPase activating protein (GAP), the protein increases the rate of conversion of the GTP to GDP. This hydrolysis allows the G alpha subunits to bind G beta/gamma subunit heterodimers, forming inactive G-protein heterotrimers, thereby terminating the signal. Alternate transcriptional splice variants of this gene have been observed but have not been thoroughly characterized.</p>
Gene ID:	10636
Pathways:	Myometrial Relaxation and Contraction , Regulation of G-Protein Coupled Receptor Protein Signaling , Platelet-derived growth Factor Receptor Signaling

Application Details

Application Notes:	WB 1:300-5000 IHC-P 1:200-400 IHC-F 1:100-500
Restrictions:	For Research Use only

Handling

Format:	Liquid
Concentration:	1 µg/µL
Buffer:	Aqueous buffered solution containing 0.01M TBS (pH 7.4) with 1 % BSA, 0.03 % Proclin300 and 50 % Glycerol.
Preservative:	ProClin
Precaution of Use:	This product contains ProClin: a POISONOUS AND HAZARDOUS SUBSTANCE, which should be handled by trained staff only.
Storage:	-20 °C
Storage Comment:	Store at -20°C for 12 months.

Handling

Expiry Date: 12 months