# antibodies -online.com





# anti-TIFA antibody (AA 51-150) (Biotin)



## Overview

Quantity:	100 μL
Target:	TIFA
Binding Specificity:	AA 51-150
Reactivity:	Mouse
Host:	Rabbit
Clonality:	Polyclonal
Conjugate:	This TIFA antibody is conjugated to Biotin
Application:	ELISA, Western Blotting (WB), Immunohistochemistry (Frozen Sections) (IHC (fro)), Immunohistochemistry (Paraffin-embedded Sections) (IHC (p))

# **Product Details**

Immunogen:	KLH conjugated synthetic peptide derived from human TIFA
Isotype:	IgG
Cross-Reactivity:	Mouse
Predicted Reactivity:	Human,Rat,Dog,Cow,Sheep,Pig,Chicken
Purification:	Purified by Protein A.

# **Target Details**

Target:	TIFA
Alternative Name:	TIFA (TIFA Products)

### **Target Details**

Background:

Synonyms: Putative MAPK activating protein PM14, Putative MAPK-activating protein PM14, Putative NF kappa B activating protein 20, Putative NF-kappa-B-activating protein 20, T2BP, TA, TA\_HUMAN, TRAF interacting protein with FHA domain containing protein A, TRAF-interacting protein with FHA domain-containing protein A, TRAF2 binding protein, TRAF2-binding protein. Background: Adapter protein which mediates the IRAK1 and TRAF6 interaction following IL-1 stimulation, resulting in the downstream activation of NF-kappa-B and AP-1 pathways. Induces the oligomerization and polyubiquitination of TRAF6, which leads to the activation of TAK1 and IKK through a proteasome-independent mechanism.

# **Application Details**

Application Notes:

WB 1:300-5000

IHC-P 1:200-400

IHC-F 1:100-500

Restrictions:

For Research Use only

# Handling

Format:	Liquid
Concentration:	1 μg/μL
Buffer:	Aqueous buffered solution containing 0.01M TBS (pH 7.4) with 1 % BSA, 0.03 % Proclin300 and 50 % Glycerol.
Preservative:	ProClin
Precaution of Use:	This product contains ProClin: a POISONOUS AND HAZARDOUS SUBSTANCE, which should be handled by trained staff only.
Storage:	-20 °C
Storage Comment:	Store at -20°C for 12 months.
Expiry Date:	12 months