

Datasheet for ABIN1700147

anti-BNIP3 antibody (pSer95) (Biotin)[Go to Product page](#)

Overview

Quantity:	100 µL
Target:	BNIP3
Binding Specificity:	pSer95
Reactivity:	Human
Host:	Rabbit
Clonality:	Polyclonal
Conjugate:	This BNIP3 antibody is conjugated to Biotin
Application:	ELISA, Immunohistochemistry (Paraffin-embedded Sections) (IHC (p)), Immunohistochemistry (Frozen Sections) (IHC (fro))

Product Details

Immunogen:	KLH conjugated synthetic phosphopeptide derived from human BNIP3 around the phosphorylation site of Ser95
Isotype:	IgG
Cross-Reactivity:	Human
Predicted Reactivity:	Dog,Cow,Sheep,Horse
Purification:	Purified by Protein A.

Target Details

Target:	BNIP3
---------	-------

Target Details

Alternative Name:	BNIP3 (BNIP3 Products)
Background:	Synonyms: BNIP3 phospho S95, p-BNIP3 phospho S95, BCL2 Adenovirus E1B 19 kDa Interacting Protein 3, BCL2/adenovirus E1B 19 kDa protein interacting protein 3, BCL2/adenovirus E1B 19 kDa protein-interacting protein 3, BNIP 3, BNIP3, BNIP3_HUMAN, NIP 3, NIP3. Background: The adenovirus E1B protein is a viral homolog of the Bcl-2 family of proteins that are involved in regulating cell death. A family of interacting proteins, which are designated Nip or Bnip and include BNIP-1, BNIP-2, BNIP-3 and Nix, associate with both the E1B protein and Bcl-2 proteins to mediate apoptotic signaling. BNIP-1 contains a hydrophobic transmembrane domain, which enables its localization to the nuclear envelope, endoplasmic reticulum and mitochondria. BNIP-2, (previously designated Nip2 and Nip21 in human and mouse respectively), shares homology with the non-catalytic domain of Cdc42 GTPase-activating protein (Cdc42GAP). Through binding to Cdc42GAP, BNIP-2 enhances the GTPase activity of Cdc42GAP, facilitating the hydrolysis of GTP bound to Cdc42 and thereby, mediating the signaling pathways involving receptor kinases, small GTPases and apoptotic proteins. Nix, which is also designated Nip3L or Bnip3L, is highly related to BNIP-3, and both proteins localize to the mitochondria where they associate with Bcl-2 proteins. BNIP-3 preferentially binds to Bcl-xL and induces apoptosis by suppressing the anti-apoptosis activity of Bcl-xL.
Gene ID:	664
Pathways:	Autophagy , Brown Fat Cell Differentiation

Application Details

Application Notes:	IHC-P 1:200-400 IHC-F 1:100-500
Restrictions:	For Research Use only

Handling

Format:	Liquid
Concentration:	1 µg/µL
Buffer:	Aqueous buffered solution containing 0.01M TBS (pH 7.4) with 1 % BSA, 0.03 % Proclin300 and 50 % Glycerol.
Preservative:	ProClin

Handling

Precaution of Use:	This product contains ProClin: a POISONOUS AND HAZARDOUS SUBSTANCE, which should be handled by trained staff only.
Storage:	-20 °C
Storage Comment:	Store at -20°C for 12 months.
Expiry Date:	12 months