

[Go to Product page](#)

Datasheet for ABIN1700247

**anti-CLEC12A antibody (AA 61-160) (Biotin)**

## Overview

Quantity:	100 µL
Target:	CLEC12A
Binding Specificity:	AA 61-160
Reactivity:	Human
Host:	Rabbit
Clonality:	Polyclonal
Conjugate:	This CLEC12A antibody is conjugated to Biotin
Application:	Western Blotting (WB), ELISA, Immunohistochemistry (Frozen Sections) (IHC (fro)), Immunohistochemistry (Paraffin-embedded Sections) (IHC (p))

## Product Details

Immunogen:	KLH conjugated synthetic peptide derived from human CLEC12A
Isotype:	IgG
Predicted Reactivity:	Human,Mouse,Rat
Purification:	Purified by Protein A.

## Target Details

Target:	CLEC12A
Alternative Name:	CLEC12A ( <a href="#">CLEC12A Products</a> )
Background:	Synonyms: C-type lectin domain family 12 member A, C-type lectin-like molecule 1,

## Target Details

CL12A\_HUMAN, CLEC12A, CLL-1, DCAL-2, Dendritic cell-associated lectin 2, MICL, Myeloid inhibitory C-type lectin-like receptor.

Background: The C-type lectin/C-type lectin-like domain (CTL/CTLD) superfamily consists of a variety of proteins that share a common protein fold and have diverse functions, including cell-cell signaling, cell adhesion, glycoprotein turnover and immune responses. CLEC-12A (C-type lectin domain family 12, member A), also known as CLL1, MICL, CLL-1 or DCAL-2, is a 275 amino acid single-pass type II membrane protein that contains one C-type lectin domain and belongs to the CTL/CTLD superfamily. Existing as multiple alternatively spliced isoforms that are expressed in neutrophils, eosinophils, monocytes and dendritic cells, CLEC-12A functions as a cell surface receptor that acts as a negative regulator of granulocyte and monocyte function and, via this activity, modulates signaling cascades. CLEC-12A is highly subject to post-translational glycosylation at its N-terminus and may also exist as a homodimer.

## Application Details

Application Notes:	WB 1:300-5000 IHC-P 1:200-400 IHC-F 1:100-500
Restrictions:	For Research Use only

## Handling

Format:	Liquid
Concentration:	1 µg/µL
Buffer:	Aqueous buffered solution containing 0.01M TBS ( pH 7.4) with 1 % BSA, 0.03 % Proclin300 and 50 % Glycerol.
Preservative:	ProClin
Precaution of Use:	This product contains ProClin: a POISONOUS AND HAZARDOUS SUBSTANCE, which should be handled by trained staff only.
Storage:	-20 °C
Storage Comment:	Store at -20°C for 12 months.
Expiry Date:	12 months