antibodies -online.com





anti-CPA3 antibody (AA 101-200) (Biotin)



Go to Product page

| \sim | | | | | |
|--------|----|----|----|-------------|---|
| () | VE | ۲۱ | /1 | \triangle | Λ |

| Quantity: | 100 μL | |
|----------------------|---|--|
| Target: | CPA3 | |
| Binding Specificity: | AA 101-200 | |
| Reactivity: | Rat | |
| Host: | Rabbit | |
| Clonality: | Polyclonal | |
| Conjugate: | This CPA3 antibody is conjugated to Biotin | |
| Application: | Western Blotting (WB), ELISA, Immunohistochemistry (Paraffin-embedded Sections) (IHC (p)), Immunohistochemistry (Frozen Sections) (IHC (fro)) | |

Product Details

| Immunogen: | KLH conjugated synthetic peptide derived from human CPA3/MC-CPA | |
|-----------------------|---|--|
| Isotype: | IgG | |
| Cross-Reactivity: | Rat | |
| Predicted Reactivity: | Human,Mouse | |
| Purification: | Purified by Protein A. | |

Target Details

| Target: | CPA3 |
|-------------------|-----------------------------|
| Alternative Name: | CPA3/MC-CPA (CPA3 Products) |

Target Details

| Background: | Synonyms: carboxypeptidase A, Carboxypeptidase A3, carboxypeptidase A3 mast cell, | |
|---------------------|---|--|
| | CBPA3_HUMAN, Cpa3, Mast cell carboxypeptidase A, mast cell carboxypeptidase A3, MC CPA, | |
| | MC-CPA. | |
| | Background: Carboxypeptidase A (CPA) is a pancreatic exopeptidase which hydrolyses the | |
| | peptide bond adjacent to the C-terminal end in polypeptide chains. Mast cell carboxypeptidase | |
| | A (MC-CPA), a part of the peptidase M14 family, is a highly conserved metalloprotease localized | |
| | to the secretory granules, along with trytases and chymases. MC-CPA is stored as an active | |
| | enzyme in the granule and is released, along with other inflammatory mediators, upon mast cell | |
| | degranulation. MC-CPA mirrors pancreatic carboxypeptidase A in cleaving COOH-terminal | |
| | aromatic and aliphatic amino acid residues. The optimum pH of MC-CPA is between neutral | |
| | and basic, depending upon the substrate. The MC-CPA gene, CPA3, resides on chromosome 3 | |
| | and contains 11 exons. | |
| Gene ID: | 1359 | |
| Pathways: | Peptide Hormone Metabolism, Regulation of Systemic Arterial Blood Pressure by Hormones | |
| Application Details | | |
| Application Notes: | WB 1:300-5000 | |
| | IHC-P 1:200-400 | |
| | IHC-F 1:100-500 | |
| Restrictions: | For Research Use only | |
| Handling | | |
| Format: | Liquid | |
| Concentration: | 1 μg/μL | |
| Buffer: | Aqueous buffered solution containing 0.01M TBS (pH 7.4) with 1 % BSA, 0.03 % Proclin300 and | |
| | 50 % Glycerol. | |
| Preservative: | ProClin | |
| Precaution of Use: | This product contains ProClin: a POISONOUS AND HAZARDOUS SUBSTANCE, which should be | |
| | handled by trained staff only. | |
| Storage: | -20 °C | |
| Storage Comment: | Store at -20°C for 12 months. | |
| | | |

| \vdash | land | lına |
|----------|-------|------|
| | iaria | шц |

Expiry Date:

12 months