antibodies - online.com







anti-QRFPR antibody (AA 151-250) (Biotin)



\sim					
()	VE	۲۱	/1	\triangle	Λ

Quantity:	100 μL
Target:	QRFPR
Binding Specificity:	AA 151-250
Reactivity:	Human
Host:	Rabbit
Clonality:	Polyclonal
Conjugate:	This QRFPR antibody is conjugated to Biotin
Application:	Western Blotting (WB), ELISA, Immunohistochemistry (Paraffin-embedded Sections) (IHC (p)), Immunohistochemistry (Frozen Sections) (IHC (fro))

Product Details

Immunogen:	KLH conjugated synthetic peptide derived from human G protein coupled receptor 103
Isotype:	IgG
Cross-Reactivity:	Human
Predicted Reactivity:	Mouse,Rat,Sheep,Horse,Rabbit
Purification:	Purified by Protein A.

Target Details

Target:	QRFPR
Alternative Name:	GPR103 (QRFPR Products)

Target Details

D = =			٠ ا ـ ـ ا ـ
Bac	KQI	ou	na:

Synonyms: AQ27, G protein coupled receptor 103, G-protein coupled receptor 103, G protein-coupled receptor 103, Orexigenic neuropeptide QRFP receptor, Pyroglutamylated RFamide peptide receptor, QRFP receptor, QRFPR, SP9155, QRFPR_HUMAN, GPCR GPR103.

Background: G protein-coupled receptors (GPRs) are a protein family of transmembrane receptors that transmit an extracellular signal (ligand binding) into an intracellular signal (G protein activation). GPR signaling is an evolutionarily ancient mechanism used by all eukaryotes to sense environmental stimuli and mediate cell-cell communication. All of the receptors have seven membrane-spanning domains and the extracellular parts of the receptor can be glycosylated. These extracellular loops also contain two highly conserved cysteine residues which create disulfide bonds to stabilize the receptor structure. GPR103 is a 455-amino acid protein with highest expression in the brain, retina, trigeminal ganglion, hypothalamus and vestibular nucleus. In peripheral tissues, GPR103 is expressed only in the heart, kidney and testis. GPR103 may regulate adrenal function. A hypothalamic neuropeptide of the RFamide family (26RFa) acts as an endogenous ligand for GPR103.

Gene ID:

84109

Application Details

Λnr	olication	MOTOC:
AIII	лисансят	140162

WB 1:300-5000

IHC-P 1:200-400

IHC-F 1:100-500

Restrictions:

For Research Use only

Handling

Format:	Liquid
Concentration:	1 μg/μL
Buffer:	Aqueous buffered solution containing 0.01M TBS (pH 7.4) with 1 % BSA, 0.03 % Proclin300 and 50 % Glycerol.
Preservative:	ProClin
Precaution of Use:	This product contains ProClin: a POISONOUS AND HAZARDOUS SUBSTANCE, which should be handled by trained staff only.
Storage:	-20 °C
Storage Comment:	Store at -20°C for 12 months.

		1.	
\vdash	land	ling	
1 1	ıaııu	11111	۰

Expiry Date:

12 months