

Datasheet for ABIN1700350

anti-IRF6 antibody (AA 1-100) (Biotin)



Overview

| Quantity: | 100 μL |
|-----------------------|---|
| Target: | IRF6 |
| Binding Specificity: | AA 1-100 |
| Reactivity: | Human, Mouse |
| Host: | Rabbit |
| Clonality: | Polyclonal |
| Conjugate: | This IRF6 antibody is conjugated to Biotin |
| Application: | Western Blotting (WB), ELISA |
| Product Details | |
| Immunogen: | KLH conjugated synthetic peptide derived from human IRF6 |
| Isotype: | IgG |
| Cross-Reactivity: | Human, Mouse |
| Predicted Reactivity: | Rat,Cow,Sheep,Pig,Rabbit |
| Purification: | Purified by Protein A. |
| Target Details | |
| Target: | IRF6 |
| Alternative Name: | IRF6 (IRF6 Products) |
| Background: | Synonyms: Interferon regulatory factor 6, IRF 6, IRF-6, IRF6, IRF6_HUMAN, LPS, OFC 6, OFC6, |

OTTHUMP00000034677, OTTHUMP00000034678, PIT, PPS, PPS1, VWS, VWS1.

Background: Interferon regulatory factor-1 (IRF-1) and IRF-2 have been identified as novel DNA-binding factors that function as regulators of both type I interferon (interferon-alpha and beta) and interferon-inducible genes. The two factors are structurally related, particularly in their N-terminal regions, which confer DNA binding specificity. In addition, both bind to the same sequence within the promoters of interferon-alpha and interferon-beta genes. IRF-1 functions as an activator of interferon transcription, while IRF-2 binds to the same cis elements and represses IRF-1 action. IRF-1 and IRF-2 have been reported to act in a mutually antagonistic manner in regulating cell growth, overexpression of the repressor IRF-2 leads to cell transformation while concomitant overexpression of IRF-1 causes reversion. IRF-1 and IRF-2 are members of a larger family of DNA binding proteins that includes IRF-3, IRF-4, IRF-5, IRF-6, IRF-7, ISGF-3 gamma p48 and IFN consensus sequence-binding protein (ICSBP).

Application Details

| Application Notes: | WB 1:300-5000 |
|--------------------|--|
| Restrictions: | For Research Use only |
| Handling | |
| Format: | Liquid |
| Concentration: | 1 μg/μL |
| Buffer: | Aqueous buffered solution containing 0.01M TBS (pH 7.4) with 1 % BSA, 0.03 % Proclin300 and 50 % Glycerol. |
| Preservative: | ProClin |
| Precaution of Use: | This product contains ProClin: a POISONOUS AND HAZARDOUS SUBSTANCE, which should be handled by trained staff only. |
| Storage: | -20 °C |
| Storage Comment: | Store at -20°C for 12 months. |
| Expiry Date: | 12 months |