antibodies -online.com





anti-RUSC1 antibody (AA 281-380) (Biotin)



Go to Product page

()	1/0	r\ /1	014	
()	ve	I V I	-v	V

Quantity:	100 μL	
Target:	RUSC1	
Binding Specificity:	AA 281-380	
Reactivity:	Human, Mouse	
Host:	Rabbit	
Clonality:	Polyclonal	
Conjugate:	This RUSC1 antibody is conjugated to Biotin	
Application:	ELISA, Western Blotting (WB), Immunohistochemistry (Frozen Sections) (IHC (fro)), Immunohistochemistry (Paraffin-embedded Sections) (IHC (p))	

Product Details

Immunogen:	KLH conjugated synthetic peptide derived from human RUSC1	
Isotype:	IgG	
Cross-Reactivity:	Human, Mouse	
Predicted Reactivity:	Rat,Cow,Sheep,Pig	
Purification:	Purified by Protein A.	

Target Details

Target:	RUSC1
Alternative Name:	RUSC1 (RUSC1 Products)

Target Details

Background:

Synonyms: DKFZp761A1822, Nesca, New molecule containing SH3 at the carboxy terminus, RUN and SH3 domain containing 1, RUN and SH3 domain containing protein 1, RUSC 1. Background: RUSC1 is a 902 amino acid protein that contains a RUN domain and a SH3 domain. RUSC1?s RUN domain is necessary for NGF induced nuclear redistribution. RUSC1 is a putative signaling adapter which may play a role in neuronal differentiation. RUSC1 seems to be involved in signaling pathways that are regulated by the prolonged activation of MAPK. RUSC2 (RUN and SH3 domain containing 2), also known as Iporin, is a 1,516 amino acid cytoplasmic protein that is widely expressed, with highest levels in brain and testis. The RUN domain of RUSC2 is required for interaction with Rab 1A, Rab 1B and GM130. It is thought that RUSC2 may possibly function as a connector between endoplasmic reticulum (ER) derived vesicle targets triggered by the Rab 1 GTPases and a signaling pathway regulated by molecules containing SH3 and/or poly-proline regions. RUSC2 also consists of a SH3 domain, suggesting a role in protein-protein interactions.

Application Details

Application Notes:	WB 1:300-5000
	IHC-P 1:200-400
	IHC-F 1:100-500
Restrictions:	For Research Use only

Handling

Format:	Liquid
Concentration:	1 μg/μL
Buffer:	Aqueous buffered solution containing 0.01M TBS (pH 7.4) with 1 % BSA, 0.03 % Proclin300 and 50 % Glycerol.
Preservative:	ProClin
Precaution of Use:	This product contains ProClin: a POISONOUS AND HAZARDOUS SUBSTANCE, which should be handled by trained staff only.
Storage:	-20 °C
Storage Comment:	Store at -20°C for 12 months.
Expiry Date:	12 months