

[Go to Product page](#)

Datasheet for ABIN1700423

anti-UCMA antibody (AA 21-120) (Biotin)

Overview

Quantity:	100 µL
Target:	UCMA
Binding Specificity:	AA 21-120
Reactivity:	Human
Host:	Rabbit
Clonality:	Polyclonal
Conjugate:	This UCMA antibody is conjugated to Biotin
Application:	Western Blotting (WB), ELISA, Immunohistochemistry (Paraffin-embedded Sections) (IHC (p)), Immunohistochemistry (Frozen Sections) (IHC (fro))

Product Details

Immunogen:	KLH conjugated synthetic peptide derived from human UCMA
Isotype:	IgG
Predicted Reactivity:	Human,Mouse,Rat
Purification:	Purified by Protein A.

Target Details

Target:	UCMA
Alternative Name:	UCMA (UCMA Products)
Background:	Synonyms: C10orf49, Gla rich protein, Gla-rich protein, Grp, GRP, OTTMUSP00000011599,

Target Details

RP23-272115.1, UCMA, Ucma-C, UCMA_HUMAN, Unique cartilage matrix associated protein, Unique cartilage matrix-associated protein C-terminal fragment, Upper zone of growth plate and cartilage matrix associated.

Background: UCMA is a 138 amino acid secreted protein that is highly expressed in resting chondrocytes in developing long bones and is thought to function in the early phase of chondrocyte differentiation. A furin-like protease processes UCMA into an N-terminal 37 amino acid peptide and a C-terminal 74 amino acid peptide, which is referred to as Unique cartilage matrix-associated protein C-terminal fragment (Ucma-C). Introduction of recombinant Ucma-C interferes with osteogenic differentiation of mesenchymal stem cells, MC3T3-E1 preosteoblasts and primary osteoblasts. This suggests that Ucma may be involved in the negative regulation of osteogenic differentiation of osteochondrogenic precursor cells at the cartilage-bone interface and in peripheral zones of fetal cartilage.

Gene ID: 221044

Application Details

Application Notes: WB 1:300-5000
IHC-P 1:200-400
IHC-F 1:100-500

Restrictions: For Research Use only

Handling

Format: Liquid

Concentration: 1 µg/µL

Buffer: Aqueous buffered solution containing 0.01M TBS (pH 7.4) with 1 % BSA, 0.03 % Proclin300 and 50 % Glycerol.

Preservative: ProClin

Precaution of Use: This product contains ProClin: a POISONOUS AND HAZARDOUS SUBSTANCE, which should be handled by trained staff only.

Storage: -20 °C

Storage Comment: Store at -20°C for 12 months.

Expiry Date: 12 months