

[Go to Product page](#)

Datasheet for ABIN1700641
anti-DGAT2 antibody (AA 251-360) (Biotin)

Overview

Quantity:	100 µL
Target:	DGAT2
Binding Specificity:	AA 251-360
Reactivity:	Human
Host:	Rabbit
Clonality:	Polyclonal
Conjugate:	This DGAT2 antibody is conjugated to Biotin
Application:	Western Blotting (WB), ELISA, Immunohistochemistry (Paraffin-embedded Sections) (IHC (p)), Immunohistochemistry (Frozen Sections) (IHC (fro))

Product Details

Immunogen:	KLH conjugated synthetic peptide derived from human DGAT2
Isotype:	IgG
Cross-Reactivity:	Human
Predicted Reactivity:	Mouse,Rat,Dog,Cow,Sheep,Pig,Chicken
Purification:	Purified by Protein A.

Target Details

Target:	DGAT2
Alternative Name:	DGAT2 (DGAT2 Products)

Target Details

Background:	<p>Synonyms: DGAT2, DGAT2_HUMAN, Diacylglycerol O acyltransferase like protein 2, Diacylglycerol O-acyltransferase 2, Diacylglycerol O-acyltransferase homolog 2 mouse, Diacylglycerol O-acyltransferase homolog 2, Diacylglycerol O-acyltransferase-like protein 2 isoform 1, Diglyceride acyltransferase 2, DKFZp686A15125, GS1999full, HMFN1045.</p> <p>Background: Glucose and insulin are anabolic signals which upregulate the transcriptions of a series of lipogenic enzymes to convert excess carbohydrate into triglycerides for efficient energy storage. Acyl-coenzyme A:diacylglycerol acyltransferase, also known as DGAT1 and ARGP1, is a microsomal enzyme that assists in the synthesis of fatty acids into triglycerides. DGAT1 catalyzes the terminal and only committed step in triacylglycerol synthesis by using diacylglycerol (DAG) and fatty acyl CoA as substrates. DGAT1 plays a fundamental role in the metabolism of cellular diacylglycerol and is important in higher eukaryotes for physiologic processes involving triacylglycerol metabolism, such as intestinal fat absorption, lipoprotein assembly, adipose tissue formation and lactation. DGAT2, which has no homology to DGAT1, differs from DGAT1 in that its activity has been shown to be inhibited by MgCl in an in vitro assay. DGAT2 is expressed primarily in liver and white adipose tissue, which suggests that it plays an important role in mammalian triglyceride metabolism.</p>
-------------	---

Gene ID:	84649
----------	-------

UniProt:	Q96PD7
----------	------------------------

Application Details

Application Notes:	WB 1:300-5000 IHC-P 1:200-400 IHC-F 1:100-500
Restrictions:	For Research Use only

Handling

Format:	Liquid
Concentration:	1 µg/µL
Buffer:	Aqueous buffered solution containing 0.01M TBS (pH 7.4) with 1 % BSA, 0.03 % Proclin300 and 50 % Glycerol.
Preservative:	ProClin
Precaution of Use:	This product contains ProClin: a POISONOUS AND HAZARDOUS SUBSTANCE, which should be

Handling

	handled by trained staff only.
Storage:	-20 °C
Storage Comment:	Store at -20°C for 12 months.
Expiry Date:	12 months