



[Go to Product page](#)

Datasheet for ABIN1700670
anti-FZR1 antibody (AA 141-240) (Biotin)

Overview

Quantity:	100 µL
Target:	FZR1
Binding Specificity:	AA 141-240
Reactivity:	Human, Mouse
Host:	Rabbit
Clonality:	Polyclonal
Conjugate:	This FZR1 antibody is conjugated to Biotin
Application:	Western Blotting (WB), ELISA, Immunohistochemistry (Paraffin-embedded Sections) (IHC (p)), Immunohistochemistry (Frozen Sections) (IHC (fro))

Product Details

Immunogen:	KLH conjugated synthetic peptide derived from human FZR1/CDC20C
Isotype:	IgG
Cross-Reactivity:	Human, Mouse
Predicted Reactivity:	Rat,Dog,Cow,Sheep,Pig,Horse,Chicken
Purification:	Purified by Protein A.

Target Details

Target:	FZR1
Alternative Name:	FZR1/CDC20C (FZR1 Products)

Target Details

Background:	<p>Synonyms: FZR, FZR2, CDC20C, Fizzy-related protein homolog, CDC20-like protein 1, Cdh1/Hct1 homolog, FZR1, FYR, KIAA1242</p> <p>Background: Key regulator of ligase activity of the anaphase promoting complex/cyclosome (APC/C), which confers substrate specificity upon the complex. Associates with the APC/C in late mitosis, in replacement of CDC20, and activates the APC/C during anaphase and telophase. The APC/C remains active in degrading substrates to ensure that positive regulators of the cell cycle do not accumulate prematurely. At the G1/S transition FZR1 is phosphorylated, leading to its dissociation from the APC/C. Following DNA damage, it is required for the G2 DNA damage checkpoint: its dephosphorylation and reassociation with the APC/C leads to the ubiquitination of PLK1, preventing entry into mitosis.</p>
Gene ID:	51343
UniProt:	Q9UM11
Pathways:	DNA Replication, Synthesis of DNA

Application Details

Application Notes:	<p>WB 1:300-5000</p> <p>IHC-P 1:200-400</p> <p>IHC-F 1:100-500</p>
Restrictions:	For Research Use only

Handling

Format:	Liquid
Concentration:	1 µg/µL
Buffer:	Aqueous buffered solution containing 0.01M TBS (pH 7.4) with 1 % BSA, 0.03 % Proclin300 and 50 % Glycerol.
Preservative:	ProClin
Precaution of Use:	This product contains ProClin: a POISONOUS AND HAZARDOUS SUBSTANCE, which should be handled by trained staff only.
Storage:	-20 °C
Storage Comment:	Store at -20°C for 12 months.
Expiry Date:	12 months