

[Go to Product page](#)

Datasheet for ABIN1700696

anti-HOOK1 antibody (AA 551-650) (Biotin)

Overview

Quantity:	100 µL
Target:	HOOK1
Binding Specificity:	AA 551-650
Reactivity:	Mouse
Host:	Rabbit
Clonality:	Polyclonal
Conjugate:	This HOOK1 antibody is conjugated to Biotin
Application:	Western Blotting (WB), ELISA, Immunohistochemistry (Paraffin-embedded Sections) (IHC (p)), Immunohistochemistry (Frozen Sections) (IHC (fro))

Product Details

Immunogen:	KLH conjugated synthetic peptide derived from human HOOK1
Isotype:	IgG
Cross-Reactivity:	Mouse
Predicted Reactivity:	Human,Rat,Dog,Cow,Sheep,Pig,Horse,Chicken,Rabbit
Purification:	Purified by Protein A.

Target Details

Target:	HOOK1
Alternative Name:	HOOK1 (HOOK1 Products)

Target Details

Background:	Synonyms: A930033L17Rik, Abnormal spermatozoon head shape, azh, h-hook1, hHK1, HK1, HOOK 1, Hook homolog 1 Drosophila, Hook1, HOOK1_HUMAN, MGC10642, OTTHUMP00000010548, OTTMUSP00000008480, Protein Hook homolog 1, RP23-80B16.4. Background: Microtubules mediate the spatial organization of diverse membrane-trafficking systems. The HOOK proteins, HOOK1, HOOK2 and HOOK3, comprise a family of cytosolic coiled-coil proteins that contain conserved N-terminal domains, which attach to microtubules, and more divergent C-terminal domains, which mediate binding to organelles. HOOK1, a cytoskeletal linker protein, may play a role in endocytic membrane trafficking. It exists as a homodimer, most likely mediated through its central coiled-coil domain. HOOK1 interacts with VPS18 and is required for spermatid differentiation, in which it is most likely involved in the positioning of the manchette microtubules and the flagellum. HOOK1 localizes primarily to the cytoplasm and does not associate with the Golgi complex, unlike HOOK3, which participates in the organization of the cis-Golgi compartment.
Gene ID:	51361
Pathways:	SARS-CoV-2 Protein Interactome

Application Details

Application Notes:	WB 1:300-5000 IHC-P 1:200-400 IHC-F 1:100-500
Restrictions:	For Research Use only

Handling

Format:	Liquid
Concentration:	1 µg/µL
Buffer:	Aqueous buffered solution containing 0.01M TBS (pH 7.4) with 1 % BSA, 0.03 % Proclin300 and 50 % Glycerol.
Preservative:	ProClin
Precaution of Use:	This product contains ProClin: a POISONOUS AND HAZARDOUS SUBSTANCE, which should be handled by trained staff only.
Storage:	-20 °C
Storage Comment:	Store at -20°C for 12 months.

Handling

Expiry Date: 12 months