antibodies -online.com





anti-PUM2 antibody (AA 901-1066) (Biotin)



()	1/0	r\ /1	014	
()	ve	I V I	-v	V

Quantity:	100 μL
Target:	PUM2
Binding Specificity:	AA 901-1066
Reactivity:	Human
Host:	Rabbit
Clonality:	Polyclonal
Conjugate:	This PUM2 antibody is conjugated to Biotin
Application:	Western Blotting (WB), ELISA, Immunohistochemistry (Paraffin-embedded Sections) (IHC (p)), Immunohistochemistry (Frozen Sections) (IHC (fro))

Product Details

Immunogen:	KLH conjugated synthetic peptide derived from human Pumilio 2
Isotype:	IgG
Predicted Reactivity:	Human, Mouse, Rat, Dog, Cow, Sheep, Pig, Horse, Chicken, Rabbit
Purification:	Purified by Protein A.

Target Details

Target:	PUM2
Alternative Name:	Pumilio 2 (PUM2 Products)
Background: Synonyms: FLJ36528, KIAA0235, MGC138251, MGC138253, PUM 2, Pum2, PUM2, PUMH2,	

Pumilio Drosphila homolog 2, Pumilio homolog 2 Drosophila, Pumilio homolog 2, Pumilio 2, PUM2_HUMAN.

Background: Pumilio 2 is a sequence-specific RNA-binding protein that regulates translation and mRNA stability by binding mRNA targets. It supports proliferation and self-renewal of stem cells by regulating the translation of key transcripts. The Pumilio gene encodes proteins that are required for development of germ stem cells in one or both sexes. The Pumilio protein interacts with the human Nanos1 protein and this interaction may play a conserved role in germ cell development. Pumilio 2 is highly expressed in testis and ovary and at lower levels in brain, heart, kidney, liver, muscle, placenta, intestine and stomach. It is also expressed in stem cells, germ cells and in most fetal tissues.

Gene ID: 23369

Pathways: Ribonucleoprotein Complex Subunit Organization

Application Details

Application Notes: WB 1:300-5000

IHC-P 1:200-400

IHC-F 1:100-500

Restrictions: For Research Use only

Handling

Format:	Liquid	
Concentration:	1 μg/μL	
Buffer:	Aqueous buffered solution containing 0.01M TBS (pH 7.4) with 1 % BSA, 0.03 % Proclin300 and 50 % Glycerol.	
Preservative:	ProClin	
Precaution of Use:	This product contains ProClin: a POISONOUS AND HAZARDOUS SUBSTANCE, which should be handled by trained staff only.	
Storage:	-20 °C	
Storage Comment:	Store at -20°C for 12 months.	
Expiry Date:	12 months	