# antibodies -online.com





# anti-FLRT1 antibody (AA 301-400) (Biotin)



Go to Product page

| $\sim$ |    |    |    |             |   |
|--------|----|----|----|-------------|---|
| ()     | VE | ۲۱ | /1 | $\triangle$ | Λ |

| Quantity:            | 100 μL  |  |
|----------------------|---|--|
| Target:              | FLRT1   |  |
| Binding Specificity: | AA 301-400  |  |
| Reactivity:          | Human   |  |
| Host:                | Rabbit  |  |
| Clonality:           | Polyclonal  |  |
| Conjugate:           | This FLRT1 antibody is conjugated to Biotin   |  |
| Application:         | Western Blotting (WB), ELISA, Immunohistochemistry (Paraffin-embedded Sections) (IHC (p)), Immunohistochemistry (Frozen Sections) (IHC (fro)) |  |

## **Product Details**

| Immunogen:            | KLH conjugated synthetic peptide derived from human FLRT1 |
|-----------------------|---|
| Isotype:              | IgG   |
| Predicted Reactivity: | Human,Mouse,Rat,Cow,Sheep,Pig,Chicken,Rabbit              |
| Purification:         | Purified by Protein A.                                    |

# **Target Details**

| Target:           | FLRT1  |  |
|-------------------|--|--|
| Alternative Name: | FLRT1 (FLRT1 Products)   |  |
| Background:       | Synonyms: Fibronectin-like domain containing leucine rich transmembrane protein 1; |  |

Fibronectin-like domain-containing leucine-rich transmembrane protein 1; FLRT1;

FLRT1\_HUMAN; Leucine rich repeat transmembrane protein FLRT1; Leucine-rich repeat transmembrane protein FLRT1.

Background: The leucine-rich (LRR) repeat is a 20-30 amino acid motif that forms a hydrophobic å/ horseshoe fold, allowing it to accommodate several leucine residues within a tightly packed core. All LRR repeats contain a variable segment and a highly conserved segment, the latter of which accounts for 11 or 12 residues of the entire LRR motif. FLRT1 (fibronectin leucine rich transmembrane protein 1) is a 646 amino acid single-pass type I membrane protein that contains one fibronectin type-III domain and ten LRR repeats. Expressed in kidney and brain, FLRT1 is thought to play a role in cell adhesion and receptor signaling. FLRT1 shares similarity with FLRT2 and FLRT3 and is subject to post-translational N-glycosylation. The gene encoding FLRT1 maps to human chromosome 11, which houses over 1,400 genes and comprises nearly 4% of the human genome.

Gene ID:

23769

### **Application Details**

Application Notes: WB 1:300-5000

IHC-P 1:200-400

IHC-F 1:100-500

Restrictions: For Research Use only

#### Handling

| Format:            | Liquid   |
|--------------------|--|
| Concentration:     | 1 μg/μL  |
| Buffer:            | Aqueous buffered solution containing 0.01M TBS (pH 7.4) with 1 % BSA, 0.03 % Proclin300 and 50 % Glycerol.         |
| Preservative:      | ProClin  |
| Precaution of Use: | This product contains ProClin: a POISONOUS AND HAZARDOUS SUBSTANCE, which should be handled by trained staff only. |
| Storage:           | -20 °C   |
| Storage Comment:   | Store at -20°C for 12 months.  |
| Expiry Date:       | 12 months  |