antibodies -online.com







anti-CRTC3 antibody (AA 151-260) (Biotin)



Quantity:	100 μL	
Target:	CRTC3	
Binding Specificity:	AA 151-260	
Reactivity:	Human	
Host:	Rabbit	
Clonality:	Polyclonal	
Conjugate:	This CRTC3 antibody is conjugated to Biotin	
Application:	Western Blotting (WB), ELISA, Immunohistochemistry (Paraffin-embedded Sections) (IHC (p)), Immunohistochemistry (Frozen Sections) (IHC (fro))	

Product Details

Immunogen:	KLH conjugated synthetic peptide derived from human CRTC3/TORC3	
Isotype:	IgG	
Predicted Reactivity:	Human,Mouse,Rat,Cow,Sheep,Horse,Rabbit	
Purification:	Purified by Protein A.	

Target Details

Target:	CRTC3	
Alternative Name:	CRTC3/TORC3 (CRTC3 Products)	
Background:	Synonyms: CREB regulated transcription coactivator 3, CREB-regulated transcription	

coactivator 3, CRTC 3, CRTC3, CRTC3_HUMAN, FLJ21868, TORC 3, TORC-3, TORC3, Transducer of CREB protein 3, Transducer of regulated cAMP response element binding protein CREB 3, Transducer of regulated cAMP response element binding protein 3, Transducer of regulated cAMP response element-binding protein 3, Transducer of regulated CREB protein 3. Background: TORC3 is a 619 amino acid protein that localizes to both the cytoplasm and the nucleus and belongs to the TORC family. Expressed in lung tissue and B and T lymphocytes, as well as in colon, brain, ovary, kidney, prostate, colon and heart, TORC3 functions as a transcriptional coactivator for CREB-1, thereby regulating the expression of CREB-activated genes, and is also thought to activate the SIK/TORC signaling pathway. TORC3 exists as multiple alternatively spliced isoforms and, in addition to its role in transcriptional activation, is thought to induce mitochondrial biogenesis, specifically in muscle cells.

Gene ID: 64784

Pathways: SARS-CoV-2 Protein Interactome, The Global Phosphorylation Landscape of SARS-CoV-2

Infection

Application Details

Application Notes: WB 1:300-5000

IHC-P 1:200-400

IHC-F 1:100-500

Restrictions: For Research Use only

Handling

Format:	Liquid
Concentration:	1 μg/μL
Buffer:	Aqueous buffered solution containing 0.01M TBS (pH 7.4) with 1 % BSA, 0.03 % Proclin300 and 50 % Glycerol.
Preservative:	ProClin
Precaution of Use:	This product contains ProClin: a POISONOUS AND HAZARDOUS SUBSTANCE, which should be handled by trained staff only.
Storage:	-20 °C
Storage Comment:	Store at -20°C for 12 months.

Н	land	lına
	iaria	шц

Expiry Date:

12 months