

[Go to Product page](#)

Datasheet for ABIN1700941
anti-FANCE antibody (AA 431-536) (Biotin)

Overview

Quantity:	100 µL
Target:	FANCE
Binding Specificity:	AA 431-536
Reactivity:	Human
Host:	Rabbit
Clonality:	Polyclonal
Conjugate:	This FANCE antibody is conjugated to Biotin
Application:	Western Blotting (WB), ELISA, Immunohistochemistry (Paraffin-embedded Sections) (IHC (p)), Immunohistochemistry (Frozen Sections) (IHC (fro))

Product Details

Immunogen:	KLH conjugated synthetic peptide derived from human FANCE
Isotype:	IgG
Predicted Reactivity:	Human,Mouse,Rat,Dog
Purification:	Purified by Protein A.

Target Details

Target:	FANCE
Alternative Name:	FANCE (FANCE Products)
Background:	Synonyms: 2810451D06Rik, AI415634, AW209126, FACE, FAE, FANCE, FANCE_HUMAN,

Target Details

Fanconi anemia complementation group E, Fanconi anemia group E protein, Protein FACE, RGD1561045.

Background: The Fanconi anemia complementation group (FANC) currently includes FANCA, FANCB, FANCC, FANCD1 (also called BRCA2), FANCD2, FANCE, FANCF, FANCG, FANCI, FANCDJ (also called BRIP1), FANCL, FANCM and FANCN (also called PALB2). The previously defined group FANCH is the same as FANCA. Fanconi anemia is a genetically heterogeneous recessive disorder characterized by cytogenetic instability, hypersensitivity to DNA crosslinking agents, increased chromosomal breakage, and defective DNA repair. The members of the Fanconi anemia complementation group do not share sequence similarity, they are related by their assembly into a common nuclear protein complex. This gene encodes the protein for complementation group E. [provided by RefSeq, Jul 2008].

Gene ID: 2178

Pathways: [DNA Damage Repair](#)

Application Details

Application Notes: WB 1:300-5000
IHC-P 1:200-400
IHC-F 1:100-500

Restrictions: For Research Use only

Handling

Format: Liquid

Concentration: 1 µg/µL

Buffer: Aqueous buffered solution containing 0.01M TBS (pH 7.4) with 1 % BSA, 0.03 % Proclin300 and 50 % Glycerol.

Preservative: ProClin

Precaution of Use: This product contains ProClin: a POISONOUS AND HAZARDOUS SUBSTANCE, which should be handled by trained staff only.

Storage: -20 °C

Storage Comment: Store at -20°C for 12 months.

Expiry Date: 12 months