

[Go to Product page](#)

Datasheet for ABIN1700945

anti-FPGT antibody (AA 251-350) (Biotin)

Overview

| | |
|----------------------|--|
| Quantity: | 100 µL |
| Target: | FPGT |
| Binding Specificity: | AA 251-350 |
| Reactivity: | Human |
| Host: | Rabbit |
| Clonality: | Polyclonal |
| Conjugate: | This FPGT antibody is conjugated to Biotin |
| Application: | Western Blotting (WB), ELISA, Immunohistochemistry (Paraffin-embedded Sections) (IHC (p)), Immunohistochemistry (Frozen Sections) (IHC (fro)) |

Product Details

| | |
|-----------------------|--|
| Immunogen: | KLH conjugated synthetic peptide derived from human FPGT |
| Isotype: | IgG |
| Predicted Reactivity: | Human,Mouse,Rat,Dog |
| Purification: | Purified by Protein A. |

Target Details

| | |
|-------------------|---|
| Target: | FPGT |
| Alternative Name: | FPGT (FPGT Products) |
| Background: | Synonyms: FPGT, FPGT_HUMAN, Fucose 1 phosphate guanylttransferase, Fucose 1 phosphate |

Target Details

guanylyltransferase, Fucose-1-phosphate guanylyltransferase, GDP beta L fucose pyrophosphorylase, GDP L fucose diphosphorylase, GDP L fucose pyrophosphorylase, GDP-L-fucose diphosphorylase, GDP-L-fucose pyrophosphorylase, GFPP.

Background: Guanylyltransferase enzymes transfer one molecule of GTP to another molecule and also function in the transfer of guanosine nucleotides to sugar molecules. The carbohydrate moieties that are generated are covalently attached to cell surfaces and are necessary to ensure a surface contour that satisfies a variety of physiological roles. L-fucose is an important sugar in complex carbohydrates that is frequently found on plant and mammalian N-linked glycans. FPGT (Fucose-1-phosphate guanylyltransferase), also known as GFPP (GDP-L-fucose pyrophosphorylase), is a 594 amino acid cytoplasmic protein that catalyzes the formation of GDP-L-fucose from L-fucose-1-phosphate and GTP. FPGT functions to reutilize the L-fucose that is produced upon glycoprotein and glycolipid turnover.

Gene ID: 8790

Application Details

Application Notes: WB 1:300-5000
IHC-P 1:200-400
IHC-F 1:100-500

Restrictions: For Research Use only

Handling

Format: Liquid

Concentration: 1 µg/µL

Buffer: Aqueous buffered solution containing 0.01M TBS (pH 7.4) with 1 % BSA, 0.03 % Proclin300 and 50 % Glycerol.

Preservative: ProClin

Precaution of Use: This product contains ProClin: a POISONOUS AND HAZARDOUS SUBSTANCE, which should be handled by trained staff only.

Storage: -20 °C

Storage Comment: Store at -20°C for 12 months.

Expiry Date: 12 months