

## Datasheet for ABIN1700975

## anti-PIK3 gamma antibody (AA 521-620) (Biotin)



Go to Product page

_					
	1//	r	Vİ	$\triangle$	۸/
	V		VI		/ V

Overview		
Quantity:	100 μL	
Target:	PIK3 gamma (PIK3CG)	
Binding Specificity:	AA 521-620	
Reactivity:	Human	
Host:	Rabbit	
Clonality:	Polyclonal	
Conjugate:	This PIK3 gamma antibody is conjugated to Biotin	
Application:	Western Blotting (WB), ELISA, Immunohistochemistry (Paraffin-embedded Sections) (IHC (p)), Immunohistochemistry (Frozen Sections) (IHC (fro))	
Product Details		
Immunogen:	KLH conjugated synthetic peptide derived from human PIK3 gamma	
Isotype:	IgG	
Cross-Reactivity:	Human	
Predicted Reactivity:	Mouse,Rat,Dog,Rabbit	
Purification:	Purified by Protein A.	
Target Details		
	PIK3 gamma (PIK3CG)	
Target:	PIK3 gamma (PIK3CG)	

## Target Details

· ·		
Background:	Synonyms: PK3CG_HUMAN, PI 3 Kinase catalytic subunit gamma, Phosphatidylinositol 4,5-	
	bisphosphate 3-kinase catalytic subunit gamma isoform, PI3-kinase subunit gamma, PI3K-	
	gamma, PI3Kgamma,	
	Background: This gene encodes a protein that belongs to the pi3/pi4-kinase family of proteins.	
	The gene product is an enzyme that phosphorylates phosphoinositides on the 3-hydroxyl group	
	of the inositol ring. It is an important modulator of extracellular signals, including those elicited	
	by E-cadherin-mediated cell-cell adhesion, which plays an important role in maintenance of the	
	structural and functional integrity of epithelia. In addition to its role in promoting assembly of	
	adherens junctions, the protein is thought to play a pivotal role in the regulation of cytotoxicity	
	in NK cells. The gene is located in a commonly deleted segment of chromosome 7 previously	
	identified in myeloid leukemias. [provided by RefSeq, Jul 2008].	
Gene ID:	5294	
UniProt:	P48736	
Pathways:	PI3K-Akt Signaling, RTK Signaling, AMPK Signaling, TLR Signaling, Inositol Metabolic Process,	
	Hepatitis C, VEGF Signaling	
Application Details		
Application Notes:	WB 1:300-5000	
	IHC-P 1:200-400	
	IHC-F 1:100-500	
Restrictions:	For Research Use only	
Handling		
Format:	Liquid	
Concentration:	1 μg/μL	
Buffer:	Aqueous buffered solution containing 0.01M TBS (pH 7.4) with 1 % BSA, 0.03 % Proclin300 and	
	50 % Glycerol.	
Preservative:	ProClin	
Precaution of Use:	This product contains ProClin: a POISONOUS AND HAZARDOUS SUBSTANCE, which should be	
	handled by trained staff only.	
Storage:	-20 °C	

## Handling

Storage Comment:	Store at -20°C for 12 months.	
Expiry Date:	12 months	