# antibodies -online.com





# anti-Glutaredoxin 2 antibody (AA 101-164) (Biotin)



( )	11/	IN	/ie	A .
	/ // <del> </del>	۱ ات	/   (−	' \/\/

Quantity:	100 μL	
Target:	Glutaredoxin 2 (GRX2)	
Binding Specificity:	AA 101-164	
Reactivity:	Human, Rat	
Host:	Rabbit	
Clonality:	Polyclonal	
Conjugate:	This Glutaredoxin 2 antibody is conjugated to Biotin	
Application:	Western Blotting (WB), ELISA, Immunohistochemistry (Paraffin-embedded Sections) (IHC (p)), Immunohistochemistry (Frozen Sections) (IHC (fro))	

#### **Product Details**

Immunogen:	KLH conjugated synthetic peptide derived from human Glutaredoxin 2	
Isotype:	IgG	
Cross-Reactivity:	Human, Rat	
Predicted Reactivity:	Mouse,Dog,Cow,Sheep,Rabbit	
Purification:	Purified by Protein A.	

## **Target Details**

Target:	Glutaredoxin 2 (GRX2)
Alternative Name:	Glutaredoxin 2 (GRX2 Products)

## **Target Details**

Background:	Synonyms: GRX2, CGI-133, Glutaredoxin-2, mitochondrial, GLRX2	
	Background: Glutathione-dependent oxidoreductase that facilitates the maintenance of	
	mitochondrial redox homeostasis upon induction of apoptosis by oxidative stress. Involved in	
	response to hydrogen peroxide and regulation of apoptosis caused by oxidative stress. Acts as	
	a very efficient catalyst of monothiol reactions because of its high affinity for protein	
	glutathione-mixed disulfides. Can receive electrons not only from glutathione (GSH), but also	
	from thioredoxin reductase supporting both monothiol and dithiol reactions. Efficiently	
	catalyzes both glutathionylation and deglutathionylation of mitochondrial complex I, which in	
	turn regulates the superoxide production by the complex. Overexpression decreases the	
	susceptibility to apoptosis and prevents loss of cardiolipin and cytochrome c release.	
Gene ID:	51022	
UniProt:	Q9NS18	
Pathways:	Cell RedoxHomeostasis	
Application Details		
Application Notes:	WB 1:300-5000	
	IHC-P 1:200-400	
	IHC-F 1:100-500	
Restrictions:	For Research Use only	
Handling		
Format:	Liquid	
Concentration:	1 μg/μL	
Buffer:	Aqueous buffered solution containing 0.01M TBS (pH 7.4) with 1 % BSA, 0.03 % Proclin300 and	
	50 % Glycerol.	
Preservative:	ProClin	
Precaution of Use:	This product contains ProClin: a POISONOUS AND HAZARDOUS SUBSTANCE, which should be	
	handled by trained staff only.	
Storage:	-20 °C	
Storage Comment:	Store at -20°C for 12 months.	
Expiry Date:	12 months	