

# Datasheet for ABIN1701103 anti-C9orf16 antibody (AA 1-83) (Biotin)

Polyclonal



Go to Product page

Quantity:	100 μL
Target:	C9orf16
Binding Specificity:	AA 1-83
Reactivity:	Human
Host:	Rabbit

Application:	ELISA, Western Blotting (WB), Immunohistochemistry (Frozen Sections) (IHC (fro)),

This C9orf16 antibody is conjugated to Biotin

Immunohistochemistry (Paraffin-embedded Sections) (IHC (p))

## **Product Details**

Overview

Clonality:

Conjugate:

Immunogen:	KLH conjugated synthetic peptide derived from human C9orf16
Isotype:	IgG
Predicted Reactivity:	Human,Mouse,Rat,Dog,Sheep,Pig,Horse
Purification:	Purified by Protein A.

### **Target Details**

Target:	C9orf16
Alternative Name:	C9orf16 (C9orf16 Products)
Background:	Synonyms: C9orf16, Chromosome 9 open reading frame 16, Cl016_HUMAN, EST00098,

FLJ12823, MGC4639, UPF0184 protein C9orf16.

Background: C9orf16 (chromosome 9 open reading frame 16) is an 83 amino acid protein that belongs to the UPF0184 (EST00098) family and is encoded by a gene that maps to human chromosome 9q34.11. Chromosome 9 consists of about 145 million bases, represents 4 % of the human genome and encodes nearly 900 genes. Thought to play a role in gender determination, deletion of the distal portion of 9p can lead to development of male to female sex reversal, the phenotype of a female with a male X,Y genotype. Hereditary hemorrhagic telangiectasia, which is characterized by harmful vascular defects, is associated with the chromosome 9 gene encoding endoglin protein, ENG. Familial dysautonomia is also associated with chromosome 9 though through the gene IKBKAP. Notably, chromosome 9 encompasses the largest interferon family gene cluster. Chromosome 9 is partnered with chromosome 22 in the translocation leading to the aberrant production of BCR-ABL fusion protein often found in leukemias.

Gene ID:

79095

#### **Application Details**

Application Notes:

WB 1:300-5000

IHC-P 1:200-400

IHC-F 1:100-500

Restrictions:

For Research Use only

#### Handling

Format:	Liquid
Concentration:	1 μg/μL
Buffer:	Aqueous buffered solution containing 0.01M TBS (pH 7.4) with 1 % BSA, 0.03 % Proclin300 and 50 % Glycerol.
Preservative:	ProClin
Precaution of Use:	This product contains ProClin: a POISONOUS AND HAZARDOUS SUBSTANCE, which should be handled by trained staff only.
Storage:	-20 °C
Storage Comment:	Store at -20°C for 12 months.
Expiry Date:	12 months