

Datasheet for ABIN1701162

anti-ATF5 antibody (AA 201-282) (Biotin)[Go to Product page](#)

Overview

Quantity:	100 µL
Target:	ATF5
Binding Specificity:	AA 201-282
Reactivity:	Mouse
Host:	Rabbit
Clonality:	Polyclonal
Conjugate:	This ATF5 antibody is conjugated to Biotin
Application:	Western Blotting (WB), ELISA

Product Details

Immunogen:	KLH conjugated synthetic peptide derived from human ATF5
Isotype:	IgG
Cross-Reactivity:	Mouse
Predicted Reactivity:	Human,Rat,Dog,Cow,Pig,Horse
Purification:	Purified by Protein A.

Target Details

Target:	ATF5
Alternative Name:	ATF5 (ATF5 Products)
Background:	Synonyms: Activating transcription factor 5 alpha/beta, Activating transcription factor 5,

Target Details

Activating transcription factor X, AFTA, ATF 5, ATF 7, ATF7, ATFX, BZIP protein ATF7, cAMP dependent transcription factor ATF 5, cAMP dependent transcription factor ATF5, Cyclic AMP dependent transcription factor ATF 5, Cyclic AMP dependent transcription factor ATF5, FLJ34666, HMFN0395, MGC102397, NAP1, NR3 associated protein, ODA 10, Transcription factor ATFX, Transcription factor like protein ODA 10, Transcription factor like protein ODA10, ATF5_HUMAN.

Background: Eukaryotic gene transcription is regulated by sequence-specific transcription factors that bind modular cis acting promoter and enhancer elements. The ATF/CREB transcription factor family binds the palindromic cAMP response element (CRE) octanucleotide TGACGTCA. The best characterized members of this gene family include CREB-1, CREB-2 (also designated ATF-4), CRE-BPa, LZIP (also designated CREB3 and Luman), CREM-1, CREM-2, ATF-1, ATF-2, ATF-3, ATF-5, ATF-6 and ATF-7. This family of proteins contain highly divergent N-terminal domains, but share a C-terminal leucine zipper for dimerization and DNA binding. ATF-5 (ATFX), which can localize to the cytoplasm or the nucleus, binds DNA as a dimer. It interacts with CCND3 and PTP4A1.

Gene ID:	22809
Pathways:	Myometrial Relaxation and Contraction

Application Details

Application Notes:	WB 1:300-5000
Restrictions:	For Research Use only

Handling

Format:	Liquid
Concentration:	1 µg/µL
Buffer:	Aqueous buffered solution containing 0.01M TBS (pH 7.4) with 1 % BSA, 0.03 % Proclin300 and 50 % Glycerol.
Preservative:	ProClin
Precaution of Use:	This product contains ProClin: a POISONOUS AND HAZARDOUS SUBSTANCE, which should be handled by trained staff only.
Storage:	-20 °C
Storage Comment:	Store at -20°C for 12 months.

Handling

Expiry Date: 12 months