antibodies -online.com





anti-DHX15 antibody (AA 101-200) (Biotin)



O -		D	4	page
1-0	$T \cap$	$Pr \cap C$	шет	nane
\cup	LO.	1 100	IUCL	Dauc

/ N	11/0	K /	\sim	A /
	ve	1 \/	ı⊢۱	Λ
\sim	' V C	1 V		v v

Quantity:	100 μL
Target:	DHX15
Binding Specificity:	AA 101-200
Reactivity:	Human
Host:	Rabbit
Clonality:	Polyclonal
Conjugate:	This DHX15 antibody is conjugated to Biotin
Application:	Western Blotting (WB), ELISA, Immunohistochemistry (Paraffin-embedded Sections) (IHC (p)), Immunohistochemistry (Frozen Sections) (IHC (fro))

Product Details

Immunogen:	KLH conjugated synthetic peptide derived from human DHX15
Isotype:	IgG
Predicted Reactivity:	Human,Mouse,Rat,Dog,Cow,Pig,Horse,Rabbit
Purification:	Purified by Protein A.

Target Details

Target:	DHX15
Alternative Name:	DHX15 (DHX15 Products)
Background:	Synonyms: ATP dependent RNA helicase #46, DBP1, DDX15, DEADH box 15, DEAH Asp Glu Ala

His box polypeptide 15, DEAH box polypeptide 15, DEAH box protein 15, HRH2, PRP43, PrPp43p, RNA, DHX15_HUMAN.

Background: DEAD-box proteins, characterized by the conserved motif Asp-Glu-Ala-Asp, are putative RNA helicases implicated in several cellular processes involving modifications of RNA secondary structure and ribosome/spliceosome assembly. Based on their distribution patterns, some members of this family may be involved in embryogenesis, spermatogenesis, and cellular growth and division. DDX15 (DEAH-box protein 15), also known as DHX15, DBP1 or HRH2, is a nuclear ATP-dependent RNA helicase that is involved in pre-mRNA splicing and is a member of the DEAH-box subfamily of DEAD-box proteins. Expressed throughout the body, DDX15 is a pre-mRNA processing factor that plays a role in spliceosome disassembly after the release of mature mRNA. When localized to the nucleoli, DDX15 is thought to interact with the La/SSB autoantigen, an RNA chaperone that functions in various intracellular processes. DDX15 is 795 amino acids in length and is the human ortholog of the S. cerevisiae protein Prp43. helicase 2.

Gene ID:

1665

Application Details

Application	Notes:

WB 1:300-5000

IHC-P 1:200-400

IHC-F 1:100-500

Restrictions:

For Research Use only

Handling

Format:	Liquid
Concentration:	1 μg/μL
Buffer:	Aqueous buffered solution containing 0.01M TBS (pH 7.4) with 1 % BSA, 0.03 % Proclin300 and 50 % Glycerol.
Preservative:	ProClin
Precaution of Use:	This product contains ProClin: a POISONOUS AND HAZARDOUS SUBSTANCE, which should be handled by trained staff only.
Storage:	-20 °C
Storage Comment:	Store at -20°C for 12 months.
Expiry Date:	12 months