

[Go to Product page](#)

Datasheet for ABIN1701245
anti-EYA2 antibody (AA 201-300) (Biotin)

Overview

Quantity:	100 µL
Target:	EYA2
Binding Specificity:	AA 201-300
Reactivity:	Rat
Host:	Rabbit
Clonality:	Polyclonal
Conjugate:	This EYA2 antibody is conjugated to Biotin
Application:	Western Blotting (WB), ELISA, Immunohistochemistry (Paraffin-embedded Sections) (IHC (p)), Immunohistochemistry (Frozen Sections) (IHC (fro))

Product Details

Immunogen:	KLH conjugated synthetic peptide derived from human EYA2
Isotype:	IgG
Cross-Reactivity:	Rat
Predicted Reactivity:	Human, Mouse, Dog, Cow, Sheep, Pig, Horse
Purification:	Purified by Protein A.

Target Details

Target:	EYA2
Alternative Name:	EYA2 (EYA2 Products)

Target Details

Background:	<p>Synonyms: EAB 1, EAB1, EYA 2, EYA2, EYA2_HUMAN, eyes absent 2 homolog Drosophila, Eyes absent homolog 2, Translation of this uORF probably lowers the translation efficiency of EYA2.</p> <p>Background: A gene on chromosome 20q13.1 encodes Eya2 (eyes absent). EYA2 is one of four members of the eyes absent family. A 271 amino acid domain at the carboxy-terminal is highly conserved amongst the members of the eyes absent family, while the PST (proline-serine-threonine)-rich amino-terminal is highly divergent. EYA2 is expressed relatively late in development in the cytoplasm of extensor tendons and ligaments of the phalangeal elements of the limb, cranial placodes, branchial arches, central nervous system, and the developing eye. Pax3 induces the expression of Eya2 in a cascade that is necessary and sufficient for myogenesis. EYA2, like EYA1, acts as a transcriptional activator in connective tissue patterning through its PST domain, which functions as a transactivation domain. EYA2 is translocated to the nucleus by Six proteins, which interact through their domain and homeodomain with EYA2. EYA2 carboxy-terminal interacts with the G Alpha z and G Alpha i 2 proteins. This interaction prevents Six proteins from translocating EYA2 to the nucleus.</p>
-------------	--

Gene ID:	2139
----------	------

Application Details

Application Notes:	WB 1:300-5000 IHC-P 1:200-400 IHC-F 1:100-500
Restrictions:	For Research Use only

Handling

Format:	Liquid
Concentration:	1 µg/µL
Buffer:	Aqueous buffered solution containing 0.01M TBS (pH 7.4) with 1 % BSA, 0.03 % Proclin300 and 50 % Glycerol.
Preservative:	ProClin
Precaution of Use:	This product contains ProClin: a POISONOUS AND HAZARDOUS SUBSTANCE, which should be handled by trained staff only.
Storage:	-20 °C
Storage Comment:	Store at -20°C for 12 months.

Handling

Expiry Date: 12 months