antibodies -online.com





anti-GALK2 antibody (AA 141-240) (Biotin)



Go to Product page

\sim					
()	VE	۲۱	/1	\triangle	Λ

Quantity:	100 μL
Target:	GALK2
Binding Specificity:	AA 141-240
Reactivity:	Rat
Host:	Rabbit
Clonality:	Polyclonal
Conjugate:	This GALK2 antibody is conjugated to Biotin
Application:	Western Blotting (WB), ELISA, Immunohistochemistry (Paraffin-embedded Sections) (IHC (p)), Immunohistochemistry (Frozen Sections) (IHC (fro))

Product Details

Immunogen:	KLH conjugated synthetic peptide derived from human GALK2/Galactokinase 2
Isotype:	IgG
Cross-Reactivity:	Rat
Predicted Reactivity:	Human,Mouse,Dog,Cow,Sheep,Pig,Rabbit
Purification:	Purified by Protein A.

Target Details

Target:	GALK2
Alternative Name:	GALK2/Galactokinase 2 (GALK2 Products)

Target Details

Background:	Synonyms: GalNAc Kinase, Galactokinase 2, GALK2, GK2, MGC1745, N-acetylgalactosamine
	kinase, GALK2_HUMAN.
	Background: GALK2 is important in the first step of the galactose metabolism pathway. GALK1,
	which belongs to the GHMP kinase family of proteins, is a crucial enzyme for galactose
	metabolism, specifically converting ?d-galactose to galactose 1-phosphate. Defects in the gene
	encoding GALK1 can cause galactosemia II, an autosomal recessive disorder characterized by
	congenital cataracts during infancy, often within the first two weeks of life. In the adult
	population, galactosemia II can cause presenile cataracts that are secondary to accumulation
	of galactitol in the lens of the eye. A second gene, GALK2, encodes an enzyme with greater
	activity against GalNAc than galactose. GALK2 has been implicated in the salvage pathway for
	the reutilization of free GalNAc derived from the degradation of complex carbohydrates.
	the reatilization of free dailyae derived from the degradation of complex earbonydrates.
Gene ID:	2585
Application Details	
Application Notes:	WB 1:300-5000
	IHC-P 1:200-400
	IHC-F 1:100-500
Restrictions:	For Research Use only
Handling	
Format:	Liquid
Concentration:	1 μg/μL
Buffer:	Aqueous buffered solution containing 0.01M TBS (pH 7.4) with 1 % BSA, 0.03 % Proclin300 and
	50 % Glycerol.
Preservative:	ProClin
Precaution of Use:	This product contains ProClin: a POISONOUS AND HAZARDOUS SUBSTANCE, which should be handled by trained staff only.
Storage:	-20 °C
Otorage.	
Storage Comment:	Store at -20°C for 12 months.
Expiry Date:	12 months