antibodies -online.com





anti-GNPDA2 antibody (AA 101-200) (Biotin)



()	11/	IN	/ie	A .
	/ // 	۱ ات	/ (−	' \/\/

Quantity:	100 μL
Target:	GNPDA2
Binding Specificity:	AA 101-200
Reactivity:	Mouse, Rat
Host:	Rabbit
Clonality:	Polyclonal
Conjugate:	This GNPDA2 antibody is conjugated to Biotin
Application:	Western Blotting (WB), ELISA, Immunohistochemistry (Paraffin-embedded Sections) (IHC (p)), Immunohistochemistry (Frozen Sections) (IHC (fro))

Product Details

Immunogen:	KLH conjugated synthetic peptide derived from human GNPDA2	
Isotype:	IgG	
Cross-Reactivity:	Mouse, Rat	
Predicted Reactivity:	Human,Dog,Cow,Sheep,Pig,Horse	
Purification:	Purified by Protein A.	

Target Details

Target:	GNPDA2
Alternative Name:	GNPDA2 (GNPDA2 Products)

Target Details

Background:	Synonyms: GlcN6P deaminase 2, Glucosamine 6 phosphate deaminase 2, Glucosamine 6
	phosphate isomerase 2, Glucosamine 6 phosphate isomerase SB52, Glucosamine-6-phosphate
	deaminase 2, Glucosamine-6-phosphate isomerase 2, Glucosamine-6-phosphate isomerase
	SB52, GNP2, GNPDA 2, Gnpda2, GNPI2_HUMAN, SB52.
	Destruction of Destination in the management of the second state of the second by the second state of the

Background: During fertilization in mammals, the sperm activates the egg by causing an increase in the level of free cytoplasmic calcium concentration. This increased calcium concentration induces a characteristic series of oscillations that trigger egg activation and early embryo development. A hamster protein named oscillin is thought to be involved in this pathway. The enzyme glucosamine-6-phosphate isomerase (GNPI) or deaminase (GNPDA1) and the related protein GNPDA2 are the human homologs of hamster oscillin. GNPDA1 and GNPDA2 catalyze the conversion of GNP to fructose-6-phosphate and ammonia. Both proteins exist as homohexamers and are ubiquitously expressed with highest expression in testis, ovary and heart. Three isoforms of GNPDA2 are expressed due to alternative splicing events.

Gene ID:

132789

Application Details

Application Notes:	WB 1:300-5000

IHC-P 1:200-400 IHC-F 1:100-500

Restrictions: For Research Use only

Handling

Format:	Liquid
Concentration:	1 μg/μL
Buffer:	Aqueous buffered solution containing 0.01M TBS (pH 7.4) with 1 % BSA, 0.03 % Proclin300 and 50 % Glycerol.
Preservative:	ProClin
Precaution of Use:	This product contains ProClin: a POISONOUS AND HAZARDOUS SUBSTANCE, which should be handled by trained staff only.
Storage:	-20 °C
Storage Comment:	Store at -20°C for 12 months.
Expiry Date:	12 months