

# Datasheet for ABIN1701333 anti-ICK antibody (AA 1-100) (Biotin)



Go to Product page

Overview	
Quantity:	100 μL
Target:	ICK
Binding Specificity:	AA 1-100
Reactivity:	Mouse, Rat
Host:	Rabbit
Clonality:	Polyclonal
Conjugate:	This ICK antibody is conjugated to Biotin
Application:	ELISA, Immunohistochemistry (Paraffin-embedded Sections) (IHC (p)), Immunohistochemistry (Frozen Sections) (IHC (fro))
Product Details	
Immunogen:	KLH conjugated synthetic peptide derived from human ICK
Isotype:	IgG
Cross-Reactivity:	Mouse, Rat
Predicted Reactivity:	Human,Dog,Cow,Pig,Horse,Rabbit
Purification:	Purified by Protein A.
Target Details	
Target:	ICK
Alternative Name:	ICK (ICK Products)

## **Target Details**

#### Background:

Synonyms: 2210420N10Rik, Al848300, ECO, ICK\_HUMAN, Heart serine threonine protein kinase, hICK, Intestinal cell MAK like kinase, Intestinal cell kinase, KIAA0936, Laryngeal cancer kinase 2, LCK2, MAK-related kinase, MAK related kinase, MGC46090, mICK, MRK, OTTMUSP00000041856, OTTMUSP00000041857, Serine/threonine-protein kinase ICK, Serine/threonine protein kinase ICK.

Background: Eukaryotic protein kinases are enzymes that belong to a very extensive family of proteins which share a conserved catalytic core common with both serine/threonine and tyrosine protein kinases. This gene encodes an intestinal serine/threonine kinase harboring a dual phosphorylation site found in mitogen-activating protein (MAP) kinases. The protein localizes to the intestinal crypt region and is thought to be important in intestinal epithelial cell proliferation and differentiation. Alternative splicing has been observed at this locus and two variants, encoding the same isoform, have been identified.

# **Application Details**

Restrictions:	For Research Use only
	IHC-F 1:100-500
Application Notes:	IHC-P 1:200-400

### Handling

Format:	Liquid
Concentration:	1 μg/μL
Buffer:	Aqueous buffered solution containing 0.01M TBS (pH 7.4) with 1 % BSA, 0.03 % Proclin300 and 50 % Glycerol.
Preservative:	ProClin
Precaution of Use:	This product contains ProClin: a POISONOUS AND HAZARDOUS SUBSTANCE, which should be handled by trained staff only.
Storage:	-20 °C
Storage Comment:	Store at -20°C for 12 months.
Expiry Date:	12 months