

[Go to Product page](#)

Datasheet for ABIN1701454
anti-PSMA4 antibody (AA 161-261) (Biotin)

Overview

Quantity:	100 µL
Target:	PSMA4
Binding Specificity:	AA 161-261
Reactivity:	Mouse, Rat
Host:	Rabbit
Clonality:	Polyclonal
Conjugate:	This PSMA4 antibody is conjugated to Biotin
Application:	Western Blotting (WB), ELISA, Immunohistochemistry (Paraffin-embedded Sections) (IHC (p)), Immunohistochemistry (Frozen Sections) (IHC (fro))

Product Details

Immunogen:	KLH conjugated synthetic peptide derived from human PSMA4
Isotype:	IgG
Cross-Reactivity:	Mouse, Rat
Predicted Reactivity:	Human,Pig,Horse,Monkey
Purification:	Purified by Protein A.

Target Details

Target:	PSMA4
Alternative Name:	PSMA4 (PSMA4 Products)

Target Details

Background:	<p>Synonyms: HC9, HsT17706, Macropain subunit C9, MGC111191, MGC12467, MGC24813, Multicatalytic endopeptidase complex subunit C9, Proteasome prosome macropain subunit alpha type 4, Proteasome alpha 4 subunit, Proteasome component C9, Proteasome subunit alpha type 4, Proteasome subunit alpha type-4, Proteasome subunit HC9, Proteasome subunit L, PSA4_HUMAN, PSC9, PSMA 4, psmA4.</p> <p>Background: The proteasome represents a large protein complex that exists inside all eukaryotes and archaea, and in some bacteria. The main function of proteasomes is to degrade unnecessary or damaged proteins by proteolysis. The most common form of the proteasome, known as the 26S Proteasome, contains one 20S Proteasome core particle structure and two 19S regulatory caps. The 20S Proteasome core is hollow and forms an enclosed cavity, where proteins are degraded, as well as openings at the two ends to allow the target protein to enter. The 20S Proteasome core particle contains many subunits, depending on the organism. All of the subunits fall into one of two types: alpha subunits, which are structural, serve as docking domains for the regulatory particles and exterior gates blocking unregulated access to the interior cavity, or beta subunits, which are predominantly catalytic. The outer two rings in the proteasome consist of seven α subunits each, and the inner two rings each consist of seven beta subunits.</p>
Gene ID:	5685
Pathways:	Mitotic G1-G1/S Phases , DNA Replication , Synthesis of DNA

Application Details

Application Notes:	WB 1:300-5000 IHC-P 1:200-400 IHC-F 1:100-500
Restrictions:	For Research Use only
Handling	
Format:	Liquid
Concentration:	1 μ g/ μ L
Buffer:	Aqueous buffered solution containing 0.01M TBS (pH 7.4) with 1 % BSA, 0.03 % Proclin300 and 50 % Glycerol.
Preservative:	ProClin

Handling

Precaution of Use:	This product contains ProClin: a POISONOUS AND HAZARDOUS SUBSTANCE, which should be handled by trained staff only.
Storage:	-20 °C
Storage Comment:	Store at -20°C for 12 months.
Expiry Date:	12 months