



[Go to Product page](#)

Datasheet for ABIN1701476  
**anti-C6orf106 antibody (AA 1-80) (Biotin)**

### Overview

Quantity:	100 µL
Target:	C6orf106
Binding Specificity:	AA 1-80
Reactivity:	Human
Host:	Rabbit
Clonality:	Polyclonal
Conjugate:	This C6orf106 antibody is conjugated to Biotin
Application:	Western Blotting (WB), Immunohistochemistry (Paraffin-embedded Sections) (IHC (p)), Immunohistochemistry (Frozen Sections) (IHC (fro)), ELISA

### Product Details

Immunogen:	KLH conjugated synthetic peptide derived from human C6orf106
Isotype:	IgG
Cross-Reactivity:	Human
Predicted Reactivity:	Mouse,Rat,Cow,Pig,Horse
Purification:	Purified by Protein A.

### Target Details

Target:	C6orf106
Alternative Name:	C6orf106 ( <a href="#">C6orf106 Products</a> )

## Target Details

---

Background:	<p>Synonyms: C6orf106, CF106_HUMAN, Chromosome 6 open reading frame 106, dJ391022.4, FLJ22195, RP3-391022.4, Uncharacterized protein C6orf106.</p> <p>Background: C6orf106 is a Making up nearly 6 % of the human genome, chromosome 6 contains around 1,200 genes within 170 million base pairs of sequence. Deletion of a portion of the q arm of chromosome 6 is associated with early onset intestinal cancer suggesting the presence of a cancer susceptibility locus. Porphyria cutanea tarda is associated with chromosome 6 through the HFE gene which, when mutated, predisposes an individual to developing this porphyria. Notably, the PARK2 gene, which is associated with Parkinson's disease, and the genes encoding the major histocompatibility complex proteins, which are key molecular components of the immune system and determine predisposition to rheumatic diseases, are also located on chromosome 6. Stickler syndrome, 21-hydroxylase deficiency and maple syrup urine disease are also associated with genes on chromosome 6. A bipolar disorder susceptibility locus has been identified on the q arm of chromosome 6. The C6orf106 gene product has been provisionally designated C6orf106 pending further characterization.</p>
-------------	--

Gene ID:	64771
----------	-------

## Application Details

---

Application Notes:	WB 1:300-5000 IHC-P 1:200-400 IHC-F 1:100-500
--------------------	---

Restrictions:	For Research Use only
---------------	-----------------------

## Handling

---

Format:	Liquid
---------	--------

Concentration:	1 µg/µL
----------------	---------

Buffer:	Aqueous buffered solution containing 0.01M TBS ( pH 7.4) with 1 % BSA, 0.03 % Proclin300 and 50 % Glycerol.
---------	---

Preservative:	ProClin
---------------	---------

Precaution of Use:	This product contains ProClin: a POISONOUS AND HAZARDOUS SUBSTANCE, which should be handled by trained staff only.
--------------------	--

Storage:	-20 °C
----------	--------

Storage Comment:	Store at -20°C for 12 months.
------------------	-------------------------------

## Handling

---

Expiry Date: 12 months