antibodies -online.com





anti-Hephaestin antibody (AA 21-120) (Biotin)



()	1 /	\sim	rv	11/	11	Α
	1//	┙	I \/	16	٦,	/\

Quantity:	100 μL	
Target:	Hephaestin (HEPH)	
Binding Specificity:	AA 21-120	
Reactivity:	Human, Mouse	
Host:	Rabbit	
Clonality:	Polyclonal	
Conjugate:	This Hephaestin antibody is conjugated to Biotin	
Application:	Western Blotting (WB), ELISA, Immunohistochemistry (Frozen Sections) (IHC (fro)), Immunohistochemistry (Paraffin-embedded Sections) (IHC (p))	

Product Details

Immunogen:	KLH conjugated synthetic peptide derived from human Hephaestin	
Isotype:	IgG	
Cross-Reactivity:	Human, Mouse	
Predicted Reactivity:	Rat	
Purification:	Purified by Protein A.	

Target Details

Target:	Hephaestin (HEPH)
Alternative Name:	Hephaestin (HEPH Products)

Target Details

Storage Comment:

Expiry Date:

rarget Details	
Background:	Synonyms: CPL, HEPH, HEPH_HUMAN, Hephaestin.
	Background: Hephaestin is a single-pass type I membrane protein that belongs to the
	multicopper oxidase family of proteins. Hephaestin, a copper-dependant ferroxidase protein, is
	crucial for iron exiting intestinal enterocytes into the circulation. It mediates the movement of
	iron across the basolateral membrane in conjunction with ferroportin 1. This is an important
	link between iron and copper metabolism in mammalian systems, as copper deficiency leads to
	reduced hephaestin and reduced iron absorption resulting in anemia. Hephaestin can bind six
	copper ions per monomer and is regulated by the homeobox transcription factor CDX2.
	Increased levels of iron leads to an increase in CDX2 expression and thus Hephaestin.
	Hephaestin is primarily detected in the intestine, but is also expressed in colon, breast, bone
	trabecural cells and fibroblasts.
Gene ID:	9843
Pathways:	Transition Metal Ion Homeostasis
Application Details	
Application Notes:	WB 1:300-5000
	IHC-P 1:200-400
	IHC-F 1:100-500
Restrictions:	For Research Use only
Handling	
Format:	Liquid
Concentration:	1 μg/μL
Buffer:	Aqueous buffered solution containing 0.01M TBS (pH 7.4) with 1 % BSA, 0.03 % Proclin300 and
	50 % Glycerol.
Preservative:	ProClin
Precaution of Use:	This product contains ProClin: a POISONOUS AND HAZARDOUS SUBSTANCE, which should be
	handled by trained staff only.
Storage:	-20 °C

Store at -20°C for 12 months.

12 months