antibodies -online.com





anti-Ulex Europaeus Lectin 1 antibody (AA 101-200) (Biotin)



Go to Product page

\sim					
()	VE	۲۱	/1	\triangle	Λ

Quantity:	100 μL
Target:	Ulex Europaeus Lectin 1
Binding Specificity:	AA 101-200
Reactivity:	Plant, Ulex europaeus (Furze)
Host:	Rabbit
Clonality:	Polyclonal
Conjugate:	This Ulex Europaeus Lectin 1 antibody is conjugated to Biotin
Application:	Western Blotting (WB), ELISA, Immunohistochemistry (Frozen Sections) (IHC (fro)), Immunohistochemistry (Paraffin-embedded Sections) (IHC (p))

Product Details

Immunogen:	KLH conjugated synthetic peptide derived from Ulex europaeus (Furze) Anti-H(0) lectin 1
Isotype:	IgG
Cross-Reactivity:	Plant
Cross-Reactivity (Details):	Ulex europaeus
Purification:	Purified by Protein A.

Target Details

Target:	Ulex Europaeus Lectin 1
Abstract:	Ulex Europaeus Lectin 1 Products

Target Details

Background:

Synonyms: UEA 1, UEA I, UEA-I, UEAI, UEA1, UEA-1, UEA1, UIEA europaeus agglutinin, Anti HO lectin 1, Anti HO lectin 1, Anti-H(0) lectin 1.

Background: Ulex Europaeus is a European gorse shrub with fragrant golden-yellow flowers.

Ulex Europaeus-I Lectin is a 46 kDa glycoprotein known to interact with -L fucosyl residues in oligosaccharides present on the membranes of human blood group O erythrocytes, human endothelial cells and a variety of human and animal epithelial cells. This antibody reacts with Ulex Europaeus Lectin 1 bound to human endothelial cells of normal and neoplastic blood and lymphatic vesicles. It also reacts with human epithelia such as in the colon, bronchus, epidermis, sweat gland ducts and hair follicles. It also reacts with for Ulex Europaeus Lectin 1 bound to vascular endothelium and squamous epithelium of human tonsil. While erythrocytes may be stained, no other tonsilar elements are reactive with the antibody. This antibody may be useful in the detection of vascular tumors, the investigation of vascular invasion by tumor cells, for determination of Ulex Europaeus Lectin 1 binding to normal, embryonal, dysplastic and neoplastic epithelial and the study of storage diseases such as fucosidosis.

UniProt:

P22972

Application Details

Δnn	lication	Notae.

WB 1:300-5000

IHC-P 1:200-400

IHC-F 1:100-500

Restrictions:

For Research Use only

Handling

Format:	Liquid
Concentration:	1 μg/μL
Buffer:	Aqueous buffered solution containing 0.01M TBS (pH 7.4) with 1 % BSA, 0.03 % Proclin300 and 50 % Glycerol.
Preservative:	ProClin
Precaution of Use:	This product contains ProClin: a POISONOUS AND HAZARDOUS SUBSTANCE, which should be handled by trained staff only.
Storage:	-20 °C
Storage Comment:	Store at -20°C for 12 months.

		1.	
\vdash	land	lın	
	ıaııu	1111	U

Expiry Date:

12 months