



[Go to Product page](#)

Datasheet for ABIN1701786  
**anti-CSTL1 antibody (AA 51-145) (Cy3)**

### Overview

|                      |   |
|----------------------|---|
| Quantity:            | 100 µL  |
| Target:              | CSTL1   |
| Binding Specificity: | AA 51-145   |
| Reactivity:          | Human   |
| Host:                | Rabbit  |
| Clonality:           | Polyclonal  |
| Conjugate:           | This CSTL1 antibody is conjugated to Cy3  |
| Application:         | Immunofluorescence (Cultured Cells) (IF (cc)), Immunofluorescence (Paraffin-embedded Sections) (IF (p)) |

### Product Details

|                       |   |
|-----------------------|---|
| Immunogen:            | KLH conjugated synthetic peptide derived from human CSTL1 |
| Isotype:              | IgG   |
| Predicted Reactivity: | Human   |
| Purification:         | Purified by Protein A.                                    |

### Target Details

|                   |   |
|-------------------|---|
| Target:           | CSTL1   |
| Alternative Name: | CSTL1 ( <a href="#">CSTL1 Products</a> )                                  |
| Background:       | Synonyms: CST1L_HUMAN, CSTL1, CTES1, Cystatin-like 1, dJ322G13.4, RCET11. |

## Target Details

---

Background: The cystatin superfamily is a well-established family of cysteine protease inhibitors. All true cystatins inhibit cysteine peptidases of the papain family, such as cathepsins, while some also inhibit legumain family enzymes. The CRES (cystatin-related epididymal spermatogenic) protein defines a new subgroup in the family 2 cystatins of the cystatin superfamily. CRES proteins lack two of the three consensus sites necessary for the cystatin inhibition of C1 cysteine proteases. They are also preferentially expressed in postmeiotic germ cells, the proximal caput epididymidis, and anterior pituitary gonadotrophs. Therefore, CRES proteins may perform unique and tissue-specific functions in the reproductive and neuroendocrine systems. As a member of the CRES subfamily, Cystatin-like 1 (CSTL1) is a 145 amino acid protein and is expressed in testis.

---

Gene ID: 128817

## Application Details

---

Application Notes: IF(IHC-P) 1:50-200  
IF(IHC-F) 1:50-200  
IF(ICC) 1:50-200

---

Restrictions: For Research Use only

## Handling

---

Format: Liquid

---

Concentration: 1 µg/µL

---

Buffer: Aqueous buffered solution containing 0.01M TBS (pH 7.4) with 1 % BSA, 0.03 % Proclin300 and 50 % Glycerol.

---

Preservative: ProClin

---

Precaution of Use: This product contains ProClin: a POISONOUS AND HAZARDOUS SUBSTANCE, which should be handled by trained staff only.

---

Storage: -20 °C

---

Storage Comment: Store at -20°C. Aliquot into multiple vials to avoid repeated freeze-thaw cycles.

---

Expiry Date: 12 months