

Datasheet for ABIN1701806

anti-FUT6 antibody (AA 101-200) (Cy3)



Go to Product page

| \sim | | | • | |
|--------|------|------|--------|----|
| |)\/A | r\ / | \cap | Λ/ |
| | | | | |

| Quantity: | 100 μL | |
|-----------------------|---|--|
| Target: | FUT6 | |
| Binding Specificity: | AA 101-200 | |
| Reactivity: | Human | |
| Host: | Rabbit | |
| Clonality: | Polyclonal | |
| Conjugate: | This FUT6 antibody is conjugated to Cy3 | |
| Application: | Immunofluorescence (Cultured Cells) (IF (cc)), Immunofluorescence (Paraffin-embedded Sections) (IF (p)) | |
| Product Details | | |
| Immunogen: | KLH conjugated synthetic peptide derived from human FUT6/FucT-VI | |
| Isotype: | IgG | |
| Predicted Reactivity: | Human | |
| Purification: | Purified by Protein A. | |
| Target Details | | |
| Target: | FUT6 | |
| Alternative Name: | FUT6 (FUT6 Products) | |
| Background: | Synonyms: Alpha 1,3 fucosyltransferase, EC=2.4.1.65, FCT3A, Fuc TVI, Fucosyltransferase 6, | |

Fucosyltransferase VI, FucT VI, Galactoside 3 L fucosyltransferase, FUT6_HUMAN. Background: Glycosyltransferases that mediate the regio- and stereoselective transfer of sugars, such as the fucosyltransferases, determine cell surface-carbohydrate profiles, which is an essential interface for biological recognition processes. Fucosyltransferases catalyze the covalent association of fucose to different positional linkages in sugar acceptor molecules. The carbohydrate moieties generated and covalently attached to cell surfaces are necessary to ensure a surface contour that satisfies physiological roles, which are reliant on adhesion molecules such as Selectins (1-3). Hematopoietic lineages rely on Fucosyltransferases to confer a surface carbohydrate phenotype, which mediates proper cell adhesion molecule recruitment and cell trafficking (4-6).

Gene ID:

2528

Application Details

Application Notes: IF(IHC-P) 1:50-200

IF(IHC-F) 1:50-200 IF(ICC) 1:50-200

Restrictions:

For Research Use only

Handling

| Format: | Liquid |
|--------------------|--|
| Concentration: | 1 μg/μL |
| Buffer: | Aqueous buffered solution containing 0.01M TBS (pH 7.4) with 1 % BSA, 0.03 % Proclin300 and 50 % Glycerol. |
| Preservative: | ProClin |
| Precaution of Use: | This product contains ProClin: a POISONOUS AND HAZARDOUS SUBSTANCE, which should be handled by trained staff only. |
| Storage: | -20 °C |
| Storage Comment: | Store at -20°C. Aliquot into multiple vials to avoid repeated freeze-thaw cycles. |
| Expiry Date: | 12 months |