# antibodies -online.com





# anti-HESX Homeobox 1 antibody (AA 51-150) (Cy3)



( )	1/0	r\ /1	014	
( )	ve	I V I	-v	V

Quantity:	100 μL	
Target:	HESX Homeobox 1 (HESX1)	
Binding Specificity:	AA 51-150	
Reactivity:	Rat	
Host:	Rabbit	
Clonality:	Polyclonal	
Conjugate:	This HESX Homeobox 1 antibody is conjugated to Cy3	
Application:	Immunofluorescence (Cultured Cells) (IF (cc)), Immunofluorescence (Paraffin-embedded Sections) (IF (p))	

### Product Details

Immunogen:	KLH conjugated synthetic peptide derived from human HESX1	
Isotype:	IgG	
Cross-Reactivity:	Rat	
Predicted Reactivity:	Human,Mouse,Dog,Cow,Sheep,Pig,Horse,Rabbit	
Purification:	Purified by Protein A.	

## **Target Details**

Target:	HESX Homeobox 1 (HESX1)
Alternative Name:	HESX1 (HESX1 Products)

#### **Target Details**

#### Background:

Synonyms: NULL

Background: The homeobox protein, HESX1, which is also known as Rathke?s pouch homeobox, HANF, homeodomain transcription factor, and anterior-restricted homeobox protein is a transcription factor that belongs to the homeodomain family of DNA binding proteins. HESX1 is initially expressed in embryonic stem cells and the primitive forebrain, and is essential for normal development of the eyes and other anterior CNS structures, such as the hypothalamus, the pituitary gland and the olfactory bulbs. The homeobox gene Hesx1 is expressed in the anterior visceral endoderm (AVE), anterior axial mesendoderm (AME), and anterior neural ectoderm (ANE) during early embryogenesis. Mutations in the Hesx1 gene are associated with disorders that are comparable with septo-optic dysplasia (SOD). These disorders are characterized by hypoplasia of the optic nerve, various types of forebrain defects and pituitary hormone deficiencies, including hypothyroidism. Hesx1 also acts as a transcriptional repressor of reporter gene constructs in tissue culture assays.

#### **Application Details**

	11	
Λnr	lication	NIOTAG
Apr	nication	110163

IF(IHC-P) 1:50-200

IF(IHC-F) 1:50-200

IF(ICC) 1:50-200

Restrictions:

For Research Use only

#### Handling

Format:	Liquid	
Concentration:	1 μg/μL	
Buffer:	Aqueous buffered solution containing 0.01M TBS (pH 7.4) with 1 % BSA, 0.03 % Proclin300 and 50 % Glycerol.	
Preservative:	ProClin	
Precaution of Use:	This product contains ProClin: a POISONOUS AND HAZARDOUS SUBSTANCE, which should be handled by trained staff only.	
Storage:	-20 °C	
Storage Comment:	Store at -20°C. Aliquot into multiple vials to avoid repeated freeze-thaw cycles.	
Expiry Date:	12 months	