# antibodies -online.com







## anti-DLG5 antibody (AA 1801-1919) (Cy3)



| ( ) | 11/               | IN   | /ie    | A .    |
|-----|-------------------|------|--------|--------|
|     | / // <del> </del> | ۱ ات | /   (− | ' \/\/ |
|     |                   |      |        |        |

| Quantity:            | 100 μL   |
|----------------------|--|
| Target:              | DLG5   |
| Binding Specificity: | AA 1801-1919   |
| Reactivity:          | Human  |
| Host:                | Rabbit   |
| Clonality:           | Polyclonal   |
| Conjugate:           | This DLG5 antibody is conjugated to Cy3  |
| Application:         | Western Blotting (WB), Immunofluorescence (Cultured Cells) (IF (cc)), Immunofluorescence (Paraffin-embedded Sections) (IF (p)) |

#### **Product Details**

| Immunogen:            | KLH conjugated synthetic peptide derived from human DLG5 |  |
|-----------------------|--|--|
| Isotype:              | IgG  |  |
| Predicted Reactivity: | Human,Mouse,Rat  |  |
| Purification:         | Purified by Protein A.                                   |  |

### **Target Details**

| Target:           | DLG5  |
|-------------------|---|
| Alternative Name: | DLG5 (DLG5 Products)  |
| Background:       | Synonyms: Discs large homolog 5 Drosophila, Discs large homolog 5, Discs large protein LP |

DLG, Discs large protein P dlg, Discs large protein P-dlg, Disks large homolog 5, DLG 5, DLG5, DLG5\_HUMAN, KIAA0583, Large type of P DLG, LP DLG, P DLG5, PDLG, Placenta and prostate DLG.

Background: Membrane-associated guanylate kinase (MAGUK) family members function as molecular scaffolds for the assembly of multiprotein complexes localizing to the plasma membrane. Several mammalian proteins related to the Drosophila tumor suppressor discslarge (dlg) gene product belong to the MAGUK family. MAGUK family members include the postsynaptic proteins PSD-93, DLG5, Pals1, PSD-95 (SAP 90), densin-180, NE-dlg (SAP 120), dlg-1 (SAP 97), GKAP (GK-associated protein), p55, the tight junction associated proteins ZO-1-3 and the caspase-associated recruitment domain (CARD) proteins CARD6, CARD8-12 and CARD14. DLG5, a cell-cell junction peripheral membrane protein, plays an important role in maintaining the structure of epithelial cell plasma membranes. It also plays an important part in transmitting extracellular signals to the cytoskeleton and the membrane. DLG5 which can interact with MPP1 and CTNNB1, is primarily expressed in prostate and placenta.

#### **Application Details**

Application Notes:

IF(IHC-P) 1:50-200

| , ipplication i totoo. | (  |  |
|------------------------|--|--|
|                        | IF(IHC-F) 1:50-200   |  |
|                        | IF(ICC) 1:50-200   |  |
| Restrictions:          | For Research Use only  |  |
| Handling               |  |  |
| Format:                | Liquid   |  |
| Concentration:         | 1 μg/μL  |  |
| Buffer:                | Aqueous buffered solution containing 0.01M TBS (pH 7.4) with 1 % BSA, 0.03 % Proclin300 and 50 % Glycerol.         |  |
| Preservative:          | ProClin  |  |
| Precaution of Use:     | This product contains ProClin: a POISONOUS AND HAZARDOUS SUBSTANCE, which should be handled by trained staff only. |  |
| Storage:               | -20 °C   |  |
| Storage Comment:       | Store at -20°C. Aliquot into multiple vials to avoid repeated freeze-thaw cycles.                                  |  |
| Expiry Date:           | 12 months  |  |