

Datasheet for ABIN1702240 anti-DCAF7 antibody (AA 2-100) (Cy3)



Go to Froduct page

Overview	
Quantity:	100 μL
Target:	DCAF7
Binding Specificity:	AA 2-100
Reactivity:	Mouse
Host:	Rabbit
Clonality:	Polyclonal
Conjugate:	This DCAF7 antibody is conjugated to Cy3
Application:	Western Blotting (WB), Immunofluorescence (Paraffin-embedded Sections) (IF (p)), Immunofluorescence (Cultured Cells) (IF (cc))
Product Details	
Immunogen:	KLH conjugated synthetic peptide derived from human WDR68
Isotype:	IgG
Cross-Reactivity:	Mouse
Predicted Reactivity:	Human,Rat,Horse
Purification:	Purified by Protein A.
Target Details	
Target:	DCAF7

Target Details

Synonyms: AN11, Dcaf7, DCAF7_HUMAN, DDB1- and CUL4-associated factor 7, HAN11,
Human anthocyanin, Petunia, Seven WD repeat protein of the AN11 family 1, SWAN 1, WD
repeat domain 68, WD repeat protein An11 homolog, WD repeat-containing protein 68, WD
repeat-containing protein An11 homolog, WDR68.
Background: Involved in craniofacial development. Acts upstream of the EDN1 pathway and is
required for formation of the upper jaw equivalent, the palatoquadrate. The activity required for
EDN1 pathway function differs between the first and second arches (By similarity). Associates
with DIAPH1 and controls GLI1 transcriptional activity. Could be involved in normal and disease
skin development. May function as a substrate receptor for CUL4-DDB1 E3 ubiquitin-protein
ligase complex.
SARS-CoV-2 Protein Interactome
IF(IHC-P) 1:50-200
IF(IHC-F) 1:50-200
IF(ICC) 1:50-200
For Research Use only
Liquid
1 μg/μL
Aqueous buffered solution containing 0.01M TBS (pH 7.4) with 1 % BSA, 0.03 % Proclin300 and
50 % Glycerol.
ProClin
This product contains ProClin: a POISONOUS AND HAZARDOUS SUBSTANCE, which should be
handled by trained staff only.
-20 °C
Store at -20°C. Aliquot into multiple vials to avoid repeated freeze-thaw cycles.