

Datasheet for ABIN1702582 anti-DAPP1 antibody (pTyr139) (Cy3)



Overview

Overview		
Quantity:	100 μL	
Target:	DAPP1	
Binding Specificity:	pTyr139	
Reactivity:	Human	
Host:	Rabbit	
Clonality:	Polyclonal	
Conjugate:	This DAPP1 antibody is conjugated to Cy3	
Application:	Western Blotting (WB), Immunofluorescence (Cultured Cells) (IF (cc)), Immunofluorescence (Paraffin-embedded Sections) (IF (p))	
Product Details		
Immunogen:	KLH conjugated synthetic phosphopeptide derived from human DAPP1 around the phosphorylation site of Tyr139	
Isotype:	IgG	
Predicted Reactivity:	Human,Mouse,Rat,Dog,Cow,Sheep,Pig,Horse,Chicken,Rabbit,Guinea Pig	
Purification:	Purified by Protein A.	
Target Details		
Target:	DAPP1	
Alternative Name:	DAPP1 (DAPP1 Products)	

Target Details

Background:	Synonyms: DAPP1 phospho Y139, p-DAPP1 phospho Y139, B cell adapter molecule of 32 kDa,		
	B lymphocyte adapter protein Bam32, B-cell adapter molecule of 32 kDa, BAM32, DAPP1,		
	DAPP1_HUMAN, DKFZp667E0716, Dual adapter for phosphotyrosine and 3-phosphotyrosine		
	and 3-phosphoinositide, Dual adaptor of phosphotyrosine and 3 phosphoinositides, hDAPP1.		
	Background: B cell adapter molecule is also designated dual adapter for phosphotyrosine and		
	3-phosphotyrosine and 3-phosphoinositide (DAPP1) or B lymphocyte adapter protein. BAM32 is		
	a B cell-associated adapter that is crucial for B cell antigen receptor signaling regulation.		
	BAM32 interacts with PtdIns and PLC g2 and, upon B cell activation, the protein is		
	phosphorylated on tyrosine residues. It is a mainly cytoplasmic protein that can translocate to		
	the cell membrane after cell stimulation. BAM32, which contains one PH domain and one SH2		
	domain, is primarily expressed in placenta and lung tissues, but can also be detected in heart,		
	liver, pancreas and brain.		
Gene ID:	27071		
Pathways:	BCR Signaling		
Application Details			
Application Notes:	IF(IHC-P) 1:50-200		
	IF(IHC-F) 1:50-200		
	IF(ICC) 1:50-200		
Restrictions:	For Research Use only		
Handling			
Format:	Liquid		
Concentration:	1 μg/μL		
Buffer:	Aqueous buffered solution containing 0.01M TBS (pH 7.4) with 1 % BSA, 0.03 % Proclin300 and		
	50 % Glycerol.		
Preservative:	ProClin		
Precaution of Use:	This product contains ProClin: a POISONOUS AND HAZARDOUS SUBSTANCE, which should be		
	handled by trained staff only.		
Storage:	-20 °C		
Storage Comment:	Store at -20°C. Aliquot into multiple vials to avoid repeated freeze-thaw cycles.		

ш	Jand	lina
г	land	1111()
•	10110	9

Expiry Date:

12 months