# antibodies -online.com





# anti-GMEB1 antibody (AA 301-400) (Cy3)



Go to Product page

$\sim$			
	N/P	r\/	i⊢₩

Quantity:	100 μL
Target:	GMEB1
Binding Specificity:	AA 301-400
Reactivity:	Human
Host:	Rabbit
Clonality:	Polyclonal
Conjugate:	This GMEB1 antibody is conjugated to Cy3
Application:	Western Blotting (WB), Immunofluorescence (Cultured Cells) (IF (cc)), Immunofluorescence (Paraffin-embedded Sections) (IF (p))

### **Product Details**

Immunogen:	KLH conjugated synthetic peptide derived from human GMEB1
Isotype:	IgG
Predicted Reactivity:	Human,Mouse,Rat,Cow,Sheep,Horse,Rabbit
Purification:	Purified by Protein A.

# **Target Details**

Target:	GMEB1	
Alternative Name:	GMEB1 (GMEB1 Products)	
Background:	Synonyms: DNA binding protein p96P, DNA-binding protein p96P, Glucocorticoid modulatory	

element binding protein 1, Glucocorticoid modulatory element-binding protein 1, GMEB 1, GMEB-1, Gmeb1, GMEB1\_HUMAN, P96P, Parvovirus initiation factor p96, P 96, P 96, P 96. Background: GMEB-1 is a 573 amino acid protein that contains one SAND domain and is a member of the KDWK family of combinatorial transcription modulators. Localized to both the cytoplasm and the nucleus, GMEB-1 forms a heterodimer with GMEB-2 (Glucocorticoid modulatory element-binding protein 2) and, once associated with GMEB-2, plays a key role in parvovirus DNA replication. In addition, GMEB-1 functions alone as a trans-acting factor that, by binding to glucocorticoid modulatory elements (GMEs) in TAT (tyrosine aminotransferase) promoters, increases intracellular sensitivity to glucocorticoid concentrations. GMEB-1 also interacts with initiator procaspases and, via this interaction, can inhibit caspase-induced apoptosis. Due to alternative splicing events, GMEB-1 is expressed as two isoforms.

Gene ID:

10691

## **Application Details**

Application Notes: IF(IHC-P) 1:50-200
IF(IHC-F) 1:50-200

IF(ICC) 1:50-200

Restrictions:

For Research Use only

#### Handling

Format:	Liquid
Concentration:	1 μg/μL
Buffer:	Aqueous buffered solution containing 0.01M TBS (pH 7.4) with 1 % BSA, 0.03 % Proclin300 and 50 % Glycerol.
Preservative:	ProClin
Precaution of Use:	This product contains ProClin: a POISONOUS AND HAZARDOUS SUBSTANCE, which should be handled by trained staff only.
Storage:	-20 °C
Storage Comment:	Store at -20°C. Aliquot into multiple vials to avoid repeated freeze-thaw cycles.
Expiry Date:	12 months