# antibodies -online.com





# anti-ZBTB43 antibody (AA 351-450) (Cy3)



Go to Product page

$\sim$			
	N/P	r\/I	i⊢₩

Quantity:	100 μL
Target:	ZBTB43
Binding Specificity:	AA 351-450
Reactivity:	Human
Host:	Rabbit
Clonality:	Polyclonal
Conjugate:	This ZBTB43 antibody is conjugated to Cy3
Application:	Western Blotting (WB), Immunofluorescence (Cultured Cells) (IF (cc)), Immunofluorescence (Paraffin-embedded Sections) (IF (p))

#### **Product Details**

Immunogen:	KLH conjugated synthetic peptide derived from human ZBTB43/ZNF297B	
Isotype:	IgG	
Predicted Reactivity:	Human,Mouse,Rat,Dog,Cow,Sheep,Horse,Rabbit	
Purification:	Purified by Protein A.	

# **Target Details**

Target:	ZBTB43	
Alternative Name:	ZBTB43/ZNF297B (ZBTB43 Products)	
Background:	Synonyms: RP11489N222, ZBT43_HUMAN, ZBTB22B, ZBTB43, Zinc Finger And BTB Domain	

Containing 43, Zinc finger and BTB domain-containing protein 22B, Zinc finger and BTB domain-containing protein 43, Zinc Finger Protein 297B, ZnF-x, ZNF297B, ZNFX.

Background: Zinc-finger proteins contain DNA-binding domains and have a wide variety of functions, most of which encompass some form of transcriptional activation or repression. The majority of zinc-finger proteins contain a Kr\_ppel-type DNA binding domain and a KRAB domain, which is thought to interact with KAP1, thereby recruiting histone modifying proteins.

Zinc finger and BTB domain-containing protein 43 (ZBTB43), also known as ZNF297B or ZBTB22B, is a 467 amino acid member of the Kr\_ppel C2H2-type zinc-finger protein family.

Localized to the nucleus, ZBTB43 contains a BTB domain, also known as a POZ domain, which inhibits DNA binding and mediates homotypic and heterotypic dimerization. Characteristics of the BTB domain and the interaction of ZBTB43 with BDP1 suggest that ZBTB43 functions as a transcription regulator.

Gene ID:

23099

## **Application Details**

Application Notes: IF(IHC-P) 1:50-200

IF(IHC-F) 1:50-200

IF(ICC) 1:50-200

Restrictions: For Research Use only

### Handling

Format:	Liquid	
Concentration:	1 μg/μL	
Buffer:	Aqueous buffered solution containing 0.01M TBS (pH 7.4) with 1 % BSA, 0.03 % Proclin300 and 50 % Glycerol.	
Preservative:	ProClin	
Precaution of Use:	This product contains ProClin: a POISONOUS AND HAZARDOUS SUBSTANCE, which should be handled by trained staff only.	
Storage:	-20 °C	
Storage Comment:	Store at -20°C. Aliquot into multiple vials to avoid repeated freeze-thaw cycles.	
Expiry Date:	12 months	