

## Datasheet for ABIN1703078 anti-RBM9 antibody (AA 65-170) (Cy3)



Of to Froduct page

()	ve	r\/i	Δ	۱۸/
$\circ$	V C	1 V		v v

Quantity:	100 μL	
Target:	RBM9	
Binding Specificity:	AA 65-170	
Reactivity:	Mouse	
Host:	Rabbit	
Clonality:	Polyclonal	
Conjugate:	This RBM9 antibody is conjugated to Cy3	
Application:	Western Blotting (WB), Immunofluorescence (Cultured Cells) (IF (cc)), Immunofluorescence	
	(Paraffin-embedded Sections) (IF (p))	

## **Product Details**

Immunogen:	KLH conjugated synthetic peptide derived from human Fox2/RBM9	
Isotype:	IgG	
Specificity:	Based on the similarity of the proteins, there is a chance that it will cross react with FOX1, but it is non-sequential homology.	
Cross-Reactivity:	Mouse	
Predicted Reactivity:	Human,Rat,Dog,Cow,Sheep,Pig,Horse,Rabbit	
Purification:	Purified by Protein A.	

## **Target Details**

Target:	RBM9	
Alternative Name:	Fox2/RBM9 (RBM9 Products)	
Background:	Synonyms: RTA, fxh, F0X2, RBM9, Fox-2, HNRBP2, HRNBP2, dJ106l20.3, RNA binding protein	
	fox-1 homolog 2, Fox-1 homolog B, Hexaribonucleotide-binding protein 2, RNA-binding motif	
	protein 9, RNA-binding protein 9, Repressor of tamoxifen transcriptional activity, RBFOX2	
	Background: RNA-binding protein that regulates alternative splicing events by binding to 5'-	
	UGCAUGU-3' elements. Prevents binding of U2AF2 to the 3'-splice site. Regulates alternative	
	splicing of tissue-specific exons and of differentially spliced exons during erythropoiesis (By	
	similarity). RNA-binding protein that seems to act as a coregulatory factor of ER-alpha.	
Gene ID:	23543	
UniProt:	043251	
Pathways:	Intracellular Steroid Hormone Receptor Signaling Pathway, Skeletal Muscle Fiber Development	
Application Details		
Application Notes:	IF(IHC-P) 1:50-200	
	IF(IHC-F) 1:50-200	
	IF(ICC) 1:50-200	
Restrictions:	For Research Use only	
Handling		
Format:	Liquid	
Concentration:	1 μg/μL	
Buffer:	Aqueous buffered solution containing 0.01M TBS (pH 7.4) with 1 % BSA, 0.03 % Proclin300 and	
	50 % Glycerol.	
Preservative:	ProClin	
Precaution of Use:	This product contains ProClin: a POISONOUS AND HAZARDOUS SUBSTANCE, which should be	
	handled by trained staff only.	
Storage:	-20 °C	
Storage Comment:	Store at -20°C. Aliquot into multiple vials to avoid repeated freeze-thaw cycles.	
Expiry Date:	12 months	