# antibodies .- online.com





## anti-GNPTAB antibody (AA 1-100) (Cy3)



Go to Product page

( )	11/0	K\ /	iew	1
	$\cup$	ועוי	$\square \vee \vee$	ı

Quantity:	100 μL	
Target:	GNPTAB	
Binding Specificity:	AA 1-100	
Reactivity:	Human	
Host:	Rabbit	
Clonality:	Polyclonal	
Conjugate:	This GNPTAB antibody is conjugated to Cy3	
Application:	Immunofluorescence (Cultured Cells) (IF (cc)), Immunofluorescence (Paraffin-embedded Sections) (IF (p))	
Product Details		
Immunogen:	KLH conjugated synthetic peptide derived from human N-acetylglucosamine-1-	
	phosphotransferase subunit alpha	
Isotype:	IgG	
Predicted Reactivity:	Human,Mouse,Rat,Cow,Pig,Horse,Rabbit	
Purification:	Purified by Protein A.	
Target Details		
	<u> </u>	
Target:	GNPTAB	

#### **Target Details**

#### Background:

Synonyms: N-acetylglucosamine-1-phosphotransferase subunit alpha, EC=2.7.8.17, GlcNAc-1-phosphotransferase subunits alpha/beta, GNPTA\_HUMAN, Gnptab, KIAA1208, Stealth protein GNPTAB, UDP-N-acetylglucosamine-1-phosphotransferase subunits alpha/beta.

Background: This gene encodes two of three subunit types of the membrane-bound enzyme N-acetylglucosamine-1-phosphotransferase, a heterohexameric complex composed of two alpha, two beta, and two gamma subunits. The encoded protein is proteolytically cleaved at the Lys928-Asp929 bond to yield mature alpha and beta polypeptides while the gamma subunits are the product of a distinct gene (GeneID 84572). In the Golgi apparatus, the heterohexameric complex catalyzes the first step in the synthesis of mannose 6-phosphate recognition markers on certain oligosaccharides of newly synthesized lysosomal enzymes. These recognition markers are essential for appropriate trafficking of lysosomal enzymes. Mutations in this gene have been associated with both mucolipidosis II and mucolipidosis IIIA.[provided by RefSeq, May 2010].

### **Application Details**

Application Notes:

**Expiry Date:** 

• •		
	IF(IHC-F) 1:50-200	
	IF(ICC) 1:50-200	
Restrictions:	For Research Use only	
Handling		
Format:	Liquid	
Concentration:	1 μg/μL	
Buffer:	Aqueous buffered solution containing 0.01M TBS (pH 7.4) with 1 % BSA, 0.03 % Proclin300 and 50 % Glycerol.	
Preservative:	ProClin	
Precaution of Use:	This product contains ProClin: a POISONOUS AND HAZARDOUS SUBSTANCE, which should be handled by trained staff only.	
Storage:	-20 °C	
Storage Comment:	Store at -20°C. Aliquot into multiple vials to avoid repeated freeze-thaw cycles.	

IF(IHC-P) 1:50-200

12 months