antibodies .- online.com





anti-Hephaestin antibody (AA 21-120) (Cy3)



Go to Product page

()	11/0	K\ /	iew	1
	\cup	'I V/I	$\square \vee \vee$	ı

Quantity:	100 μL
Target:	Hephaestin (HEPH)
Binding Specificity:	AA 21-120
Reactivity:	Human, Mouse
Host:	Rabbit
Clonality:	Polyclonal
Conjugate:	This Hephaestin antibody is conjugated to Cy3
Application:	Western Blotting (WB), Immunofluorescence (Cultured Cells) (IF (cc)), Immunofluorescence (Paraffin-embedded Sections) (IF (p))

Product Details

Immunogen:	KLH conjugated synthetic peptide derived from human Hephaestin
Isotype:	IgG
Cross-Reactivity:	Human, Mouse
Predicted Reactivity:	Rat
Purification:	Purified by Protein A.

Target Details

Target:	Hephaestin (HEPH)
Alternative Name:	Hephaestin (HEPH Products)

Target Details

Background:	Synonyms: CPL, HEPH, HEPH_HUMAN, Hephaestin.	
	Background: Hephaestin is a single-pass type I membrane protein that belongs to the	
	multicopper oxidase family of proteins. Hephaestin, a copper-dependant ferroxidase protein, is	
	crucial for iron exiting intestinal enterocytes into the circulation. It mediates the movement of	
	iron across the basolateral membrane in conjunction with ferroportin 1. This is an important	
	link between iron and copper metabolism in mammalian systems, as copper deficiency leads to	
	reduced hephaestin and reduced iron absorption resulting in anemia. Hephaestin can bind six	
	copper ions per monomer and is regulated by the homeobox transcription factor CDX2.	
	Increased levels of iron leads to an increase in CDX2 expression and thus Hephaestin.	
	Hephaestin is primarily detected in the intestine, but is also expressed in colon, breast, bone	
	trabecural cells and fibroblasts.	
Gene ID:	9843	
Pathways:	Transition Metal Ion Homeostasis	
Application Details		
Application Notes:	IF(IHC-P) 1:50-200	
	IF(IHC-F) 1:50-200	
	IF(ICC) 1:50-200	
Restrictions:	For Research Use only	
Handling		
Format:	Liquid	
Concentration:	1 μg/μL	
Buffer:	Aqueous buffered solution containing 0.01M TBS (pH 7.4) with 1 % BSA, 0.03 % Proclin300 and	
	50 % Glycerol.	
Preservative:	ProClin	
Precaution of Use:	This product contains ProClin: a POISONOUS AND HAZARDOUS SUBSTANCE, which should be	
	handled by trained staff only.	
Storage:	-20 °C	
Storage Comment:	Store at -20°C. Aliquot into multiple vials to avoid repeated freeze-thaw cycles.	
Expiry Date:	12 months	